

# **The potential effects of microplastics on human health – what is known and what is unknown**

## **Abstract**

Microplastic contamination is ubiquitous in aquatic and terrestrial environments, found in water, sediments, within organisms and in the atmosphere and the biological effects on animal and plant life have been extensively investigated in recent years. There is growing evidence that humans are exposed to microplastics via ingestion of food and drink and through inhalation. Despite the prevalence of contamination, there has been limited research on the effects of microplastics on human health and most studies, to date, analyse the effects on model organisms with the likely impacts on human health being inferred by extrapolation. This review summarises the latest findings in the field with respect to the prevalence of microplastics in the human environment, to what extent they might enter and persist in the body, and what effect, if any, they are likely to have on human health. While definitive evidence linking microplastic consumption to human health is currently lacking, results from correlative studies in people exposed to high concentrations of microplastics, model animal and cell culture experiments, suggest that effects of microplastics could include provoking immune and stress responses and inducing reproductive and developmental toxicity. Further research is required to explore the potential implications of this recent contaminant in our environment in more rigorous clinical studies.

## **Keywords**

Microplastics, human health risks, POPs, inhalation, ingestion

## Introduction

Large scale production of plastics dates back to around the 1950s (Boucher and Billard 2019). In 2010, 275 million metric tons (MT) of plastic waste was generated in 192 coastal countries (Jambeck et al. 2015). By 2017 this rose to 335 million MT of plastic waste (Boucher and Billard 2019). It is also estimated that between 4.8 and 12.7 million MT of this waste enters the world's oceans (Jambeck et al. 2015). The impact of macroplastics (> 5mm in at least one dimension) through entanglement, choking and strangulation on animals has been well documented and is the more conspicuous plastic waste that is often seen in photographs depicting the scale of the plastic waste problem. Although not as obvious to the naked eye, smaller pieces of plastic debris deemed "microplastics" (MPs) (particles < 5mm in diameter but larger than 1µm (Hartmann et al. 2019) are the most abundant form of solid waste on Earth. MPs can be further categorized into primary and secondary. Primary MPs are originally made to be micro-sized, usually for use in cosmetic products such as microbeads (Hartmann et al. 2019). Secondary MPs refers to those that have been broken down by photo degradation or mechanical weathering over time and now fall into the <5mm definition.

MPs have been found in the ocean (Law and Thompson 2014; Auta, Emenike, and Fauziah 2017; Boucher and Friot 2017), in freshwater, (Horton et al. 2017; Vaughan, Turner, and Rose 2017; Li, Liu, and Chen 2018), in sediments (Abidli et al. 2018; Reed et al. 2018), in soils (Watteau et al. 2018; Zhang et al. 2018) and in the air (Prata 2018; Wright et al. 2020). Some of these locations are quite remote, far from human settlements. MPs have been found in remote polar regions, specifically with high concentrations seen in sea ice cores (Peeken et al. 2018). Most of the MPs detected in these ice cores were smaller than 50µm and an average of 67% of the particles were within the smallest detectable class size of 11µm (Peeken et al. 2018). They are also found in the deepest parts of the world's oceans; the

Mariana Trench sediment was found to have between 200 and 2200 pieces per litre, with the majority being plastic microfibrils measuring 1-3mm in length in seawater and 0.1-0.5mm in length in sediment (Peng et al. 2018). MPs appearing in remote locations can be explained by the plastic cycle (Horton and Dixon 2018), whereby MPs accumulated in the world's oceans are so small that they can be present in the evaporation that forms our rain clouds, this rainfall containing MPs is then deposited in mountainous regions and other remote locations. The subsequent lakes and rivers transport the MPs back to the ocean, forming the plastic cycle (Geyer, Jambeck, and Law 2017; Bank and Hansson 2019). China's largest inland lake – Qinghai Lake – was found to have MPs present, with small MPs (0.1 – 0.5mm) mostly on the surface water and larger MPs (1 – 5mm) were more abundant in the connected river samples (Xiong et al. 2018). Surface water and sediment samples were collected from 6 sites along 5 different rivers in the Tibet Plateau (Jiang et al. 2019). The surface water had 483 – 967 items m<sup>-3</sup> and the sediment 50 – 195 items kg<sup>-1</sup>. These examples emphasise how widespread MP contamination has become.

We know that MPs are prevalent in oceans, lakes and rivers but are humans exposed to them? A review of MPs in commercial salt for human consumption found that in 128 brands of salt from 38 countries contained MPs (Peixoto et al. 2019). Similarly, MPs in bottled drinking water was found to be from the caps and could also have long-term exposure implications (Choudhary, Kurien, and Srivastava 2020). MPs have also been found in beer, energy drinks and other soft drinks (Kosuth, Mason, and Wattenberg 2018; Shruti et al. 2020) and more recently MPs (<10µm in diameter) have been found within the flesh of fruit and vegetables (Oliveri Conti et al. 2020).

There are many studies on how these MPs are ingested or inhaled and the effects this might have on wildlife (Lehner et al. 2019; Prata et al. 2020). The question is, to what extent are humans exposed and how could it affect humans? Despite the lack of knowledge about direct impacts of human health, it is acknowledged that plastic and micro plastic debris needs to be addressed (Katyal, Kong, and Villanueva 2020). Literature on the effects of MPs on other wildlife can be used as an indication of how they may impact human health and are summarised in this review. The effects of MPs on human health can be separated into three main categories; chemical, physical and biological effects and then further divided by exposure route and the potential clinical effects, as illustrated by the diagram in Figure 1.

## **Chemical Effects**

### Toxic additives

There is evidence to suggest that additives such as dyes or plasticisers could cause toxicity, carcinogenicity and mutagenicity (Gasperi et al. 2018). Additives, dyes and pigments could leach from MPs and accumulate on surfaces and in water sources, with the health consequences of this unknown (Gasperi et al. 2018).

Phthalates are commonly used as plasticizers to provide flexibility to plastics. They are an additive, therefore not chemically bound (covalently bonded) to the polymer and so are more likely to be released and transfer to the environment. Over 80% of plasticizers used worldwide are phthalates. They have been shown to appear in household dust (Abb et al. 2009; Ait Bamai, et al. 2014a), human urine (Jornet-Martínez, Antón-Soriano, and Campíns-Falcó 2015) and breastmilk (Main et al. 2006; Högberg et al. 2008). There is some evidence to suggest an association between the level of phthalates and occurrence of asthma and allergies, especially in children (Ait Bamai, et al. 2014b). Exposure to phthalates has also

100 been shown to have a biological effect in utero and could be associated with a shorter  
101 pregnancy duration (Latini et al. 2003). Bisphenol-A (BPA) has also been studied similarly to  
102 phthalates and shown to be a reproductive toxicant, being associated with adverse birth  
103 outcomes (Peretz et al. 2014). Monitoring of human tissues and body fluids allows us to see  
104 what concentrations of environmental contaminants are present. Biomonitoring has shown  
105 that chemicals used in the manufacture of plastics, such as BPA, phthalates and styrene, are  
106 present in the human population (Galloway 2015). Some of these chemicals have a  
107 widespread presence in the general population at concentrations capable of causing harm in  
108 animal models which raises a public health concern (Talsness et al. 2009).

109  
110 Data from a study on short-tailed shearwater birds suggested that there was a transfer of  
111 plastic derived chemicals from ingested plastics to the tissues of the birds (Tanaka et al.  
112 2013). They found brominated chemicals that were not present in the natural prey of the bird  
113 but likely from the plastic that was also found in the stomachs of some of the birds.

114  
115 In a study assessing hazard levels, 31 out of 55 polymers were composed of monomers that  
116 were assigned to the most severe hazard levels (Lithner, Larsson, and Dave 2011). Polyvinyl  
117 chloride (PVC) has a carcinogenic monomer and several hazardous additives making it  
118 arguably the most dangerous plastic in terms of toxicity. An investigation into whether  
119 various plastic products emitted hazardous chemical substances into water containing  
120 *Daphnia magna* found that 9 of the 32 products caused acute toxic effects (immobility)  
121 (Lithner et al. 2009). It was also found that PVC and polyurethane leachates were the only  
122 plastic types tested that displayed toxicity.

There have been a few studies on how MPs and their additives can cause toxic effect at a cellular level, looking at cytotoxicity, oxidative stress and cell viability. Human cerebral and epithelial cells were exposed to different levels of contaminants and showed oxidative stress when MPs were introduced but there was no significant reduction in cell viability (Schirizzi et al. 2017). Another study found that the positive control induced a high degree of toxicity in all in vitro tests using direct contact (Van Tienhoven et al. 2006). An additional study found that direct contact of polypropylene MPs with human cells could induce productions of cytokines and histamines (Hwang et al. 2019).

Polybrominated diphenyl ethers (PBDEs) are additives used as flame retardants in many commercial products. Plastic can integrate up to 15% PBDEs and they are not chemically bound so are likely to leach during production, disposal and recycling processes (Domingo 2012). Concentrations of PBDEs have increased over the years in the bodies of wildlife and humans, with the long lasting effects unknown (Linares, Bellés, and Domingo 2015). There has been a handful of studies that have indicated that bioaccumulation could cause impaired neurological development (Bellés et al. 2010; Reverte, Domingo, and Colomina 2014) and endocrine disruption (Alonso et al. 2010), however, all studies were conducted on mice or rats, therefore no conclusions can be drawn on the danger to humans.

#### Secondary toxins

The interaction of MPs and chemical pollutants is an area widely studied (Crawford and Quinn 2017). Persistent organic pollutants (POPs) are extensively recognised to be throughout the environment, including Oceans. These pollutants are hydrophobic and have been found to readily adsorb to MPs (Velzeboer, Kwadijk, and Koelmans 2014). There are many examples of this interaction in the literature, (Rios et al. 2010; Zarfl and Matthies 2010;

Frias, Sobral, and Ferreira 2010; Bakir, Rowland, and Thompson 2012; Driedger et al. 2015) as well as the MP interaction with heavy metals, (Ashton, Holmes, and Turner 2010; Holmes, Turner, and Thompson 2012; Holmes and Thompson 2014; Rochman, Hentschel, and Teh 2014). For example, polyethylene mulching sheets, used in agriculture, easily fragment into MPs. The longer the plastic mulching sheets are used in agriculture, the more microplastics that can be found in the soil indicating that they are a major source of MPs into arable soil (Huang et al. 2020a). They can also adsorb pesticides that are either in the soil or already sprayed onto the plastic (Wang et al. 2020). Carbendazim, dipterex and malathion are examples of the pesticides that can adsorb to MPs. It is suggested that MPs could become the source or carrier of pesticides into other environments, such as water, and have the potential for environmental and human safety risks. A study showed that by increasing the MP dosage there was increased removal of pesticides from solution reaching equilibrium in 120 minutes (Wang et al. 2020). They concluded that the adsorption was a spontaneous and exothermic process. In this way, additive or sorbed chemicals including polycyclic aromatic hydrocarbons (PAHs), antimicrobials, and halogenated flame retardants (HFRs) have been found in laboratory studies to be transferred from MPs to marine organisms (Browne et al. 2013; Avio et al. 2015), however the accumulation of chemical burdens from ingested MPs is not always unidirectional and depends on the concentration gradient between the ingested plastics and the gut of the organism (Koelmans et al. 2016; Bakir et al. 2016). For example, if an animal which already has a high concentration of chemical contamination from the environment ingests microplastics with low concentrations of chemicals, the transfer is expected to be from the gut to the microplastics, essentially “cleaning out” the animal.

Additionally, antibiotics can adsorb to MPs in contaminated waters and result in them being transported long distances. A study found that polyamide (PA) had the highest adsorption of

antibiotics (Li, Zhang, and Zhang 2018). Five antibiotics (sulfadiazine, amoxicillin, tetracycline, ciprofloxacin and trimethoprim) and five MPs (polyethylene, polystyrene, polypropylene, PA and PVC) were investigated in both freshwater and seawater, with a higher adsorption rate of antibiotics found in freshwater compared to seawater. The same adsorption kinetics have been studied with steroid hormones, although in less depth. It has been found that  $17\beta$ -estradiol and  $17\alpha$ -ethynylestradiol, types of synthetic hormone, will readily adsorb to MPs (Lu et al. 2020).

The potential ingestion of MPs and subsequent pollutants poses a toxic problem for the food web. Some studies have looked at how these pollutants might affect organisms (Besseling et al. 2013; Browne et al. 2013) and also look at the bioaccumulation of MPs and the pollutants (Koelmans et al. 2013). One study collected edible oysters from a coastal city in China and found MPs in all of the oyster tissue samples (Zhu et al. 2020). In addition, there was bioaccumulation of trace metals in higher concentrations than normal in the oyster tissue. It was concluded that this could pose a potential danger to humans if marine life were exposed to MPs and contaminants and then consumed.

MPs and POPs have also been found to lead to immunotoxicity in blood clams, however the larger MP size of  $30\mu\text{m}$  compared to the smaller  $500\text{nm}$  diameter appeared to mitigate the toxicity (Tang et al. 2020). On the other hand, one study concluded that the importance of MPs being a vector of toxic substances to marine organisms was of limited importance, in relation to other exposure pathways (Gouin et al. 2011). Additionally, a study on *Talitrus saltatory*, a type of amphipod, demonstrated that ingestion of contaminated MPs (polybrominated diphenyl ether) transferred organic pollutants to its tissues. However, uncontaminated MPs ingested by a contaminated amphipod removed the organic pollutants



instead (Scopetani et al. 2018). This two-way transfer could support the view that toxic substance adsorption to MPs has an equilibrium effect on marine life and is therefore a less important pathway. More research should be carried out to establish the potential of exposure to toxic pollutants carried by MPs and how they might affect human health (Rodrigues et al. 2019).

## **Physical Effects**

### **Inhalation**

There have been many studies that show there are MPs in the atmosphere that can be readily inhaled (Liu, et al. 2019b; Zhang et al. 2020b; Huang et al. 2020b). Production of plastic textile fibres has increased more than 6% per year and makes up around 16% of the worlds plastic production (Gasperi et al. 2018). Small fibres can shed from clothing due to general wear and washing, with just one garment predicted to release 1900 fibres per wash into waste water (Browne et al. 2011). The scale of plastic fibre production worldwide and the subsequent potential to be inhaled or ingested suggests investigations on their effects to human health should be considered.

There is research that has been conducted and is continuing to be produced, that is estimating the volume of airborne MPs across the globe. A study conducted in Central London tested atmospheric MP deposition and found it was 20 times greater than in a more remote location (Wright et al. 2020). They also found that fibrous MPs made up the vast majority of the plastics found (92%). Suspended atmospheric MPs were tested for in Shanghai, finding 0 – 4.18 m<sup>-3</sup> (items per cubic meter of air) (Liu et al. 2019a). Of these 67% were microfibres, 30% fragments and 3% granules, leading to the assumption that the likely source of the majority of MPs were synthetic textiles. They also estimated that people in Shanghai inhaled

approximately 21 MP particles per day whilst outdoors. An earlier study of atmospheric fallout in Dongguan City also found the dominant MP type to be fibres (Cai et al. 2017). Similar quantities of MPs were sampled over the West Pacific Ocean, suspended in marine air, with 60% microfibrils, 31% fragments and 8% granules (Liu et al. 2019c). This supports the conclusion that the vast majority of atmospheric MPs comes from synthetic textiles. It was found that daytime collection had twice the amount as the night time collection. MPs have also been found in dust in Tehran with 88-605 MP particles per 30g of dry dust (Dehghani, Moore, and Akhbarizadeh 2017). They also estimate that outdoor activity can lead to an estimated exposure of 3223 particles per year for children and 1063 particles per year for adults. A similar study in Iran found on average 900 MP particles in 15g of street dust (Abbasi et al. 2019).

Atmospheric MPs can also be sourced from deposition, or rain. A study in Paris detected MPs in atmospheric fallout, with the results finding 29 – 280 particles m<sup>-2</sup> day<sup>-1</sup> (Dris et al. 2015). Atmospheric deposition has also been tested in remote environments (Zhang et al. 2019). One study found 249 fragments, 73 films and 44 fibres per square meter in the catchment area of the French Pyrenees (Allen et al. 2019). They concluded that the MPs could travel up to 95km to reach more remote areas via atmospheric transport. A similar study instead looked at a glacier in the Italian Alps and found 74.4 MP items kg<sup>-1</sup> of sediment (Ambrosini et al. 2019). This contained most commonly polyesters, but also polyamide, polyethylene and polypropylene. Furthermore, they estimate that the whole glacier could have 131-162 million plastic items.

The effects on human health of inhaling these fibrous MPs is little understood. It is thought that the majority of fibres can be cleared from the respiratory system, however some will go

on to cause inflammatory responses and even respiratory lesions (Prata 2018), especially in those with compromised clearance mechanisms (Gasperi et al. 2018). Of 114 lung specimens from patients undergoing lung resection for removal of a tumour, 87% were observed to contain cellulosic or plastic fibres, demonstrating that these small fibres are respirable and accumulate in lung tissue (Pauly et al. 1998). Synthetic textiles are thought to be the main source of airborne MPs, especially indoors where the concentration is greater (Dris et al. 2017). It has been studied previously that inhalation of fibres during factory work can cause some cancers (Gallagher et al. 2015), however some studies on nylon flock workers suggested that there was no evidence of an increased cancer risk, but there was a higher prevalence of respiratory irritation (Wright and Kelly 2017).

We know that the lungs are exposed daily to pollutants which act as oxidants and this leads to oxidative stress, inflammation and carcinogenesis (Valavanidis et al. 2013). There is also an association between the increased incidence of respiratory disease and lung cancers from the exposure to low levels of respirable fibres (Valavanidis et al. 2013). However, little research has been conducted on the potential adverse health effects on human lungs when inhaling synthetic fibres and is therefore difficult to attribute this increase in respiratory disease to inhaled MPs (Gasperi et al. 2018). A study investigating proinflammatory responses in rats to various sizes of polystyrene particles found that the smaller particles (64nm) gave a significantly greater neutrophil influx in the lungs compared to the larger particles (202nm – 535nm) (Brown et al. 2001). It is thought that this is due to the larger surface area of smaller particles, leading to increased inflammation. There is evidence to suggest that MPs could also translocate to other tissues once inhaled or ingested, with one study finding that fluorescent polystyrene microspheres delivered intranasally to mice could be found in the spleen 10 days later (Eyles et al. 2001). It was also found that once there, they could incite immunological

functions. There is little to no information available on any human studies looking at health effects of MP fibre or particle inhalation, something that should be investigated in the future.

### Ingestion

A considerable amount of MP research is conducted on ingestion by aquatic life, (Possatto et al. 2011; Lusher, McHugh, and Thompson 2013; Phillips and Bonner 2015; Romeo et al. 2015; Barboza et al. 2018) on seabirds, (van Franeker et al. 2011; Lavers, Bond, and Hutton 2014; Amélineau et al. 2016) and other wildlife (Huerta Lwanga et al. 2016), with limited studies conducted on human ingestion (Ribeiro et al. 2019). One study compared human ingestion of MPs from mussels with the inhalation of microfibres whilst eating that same meal and found that you inhale more synthetic microfibres sitting down for a meal than you would from eating the mussels (Catarino et al. 2018). There is however, some proof that humans ingest MPs, when one study tested 8 human stool samples and found MPs in all of them (Schwabl et al. 2019). They found that polypropylene and polyethylene terephthalate were the most abundant types of plastic. It is also known that MPs are present in seafood (Li et al. 2016), water, salt and beer (Kosuth, Mason, and Wattenberg 2018) and in potentially more food or drink items that are regularly consumed by humans (Zhang et al. 2020a).

An evaluation of the number of MPs consumed from the average intake of food found that the average annual consumption was in the range of 39 000 – 52 000 particles (Cox et al. 2019). This could increase to between 74 000 – 121 000 when the inhalation of MPs was considered. They also found that if you included water intake from only a bottle source then an individual could be ingesting a further 90 000 particles compared to 40 000 particles consumed from tap water. Edible fruits and vegetables provide further example of the potential ingestion of MPs by humans. A study found that apples were the fruit most

contaminated with a median of 223 000 p/g of MPs. It was further calculated that the estimated daily intake of MPs from apples was 4.62 E+05 for adults and 1.41E+05 for children (Oliveri Conti et al. 2020).

One study in China comparing MPs in fish and bivalves in cities with other countries in the world, found that MPs are prevalent in commercial fish and bivalves sold in city markets and that the risk to human health is greater from these markets than other countries in the world (Fang et al. 2019). A more recent study looked at 150 fish from 3 species and found 49% had MPs, of this 32% was found in the dorsal muscle of the fish with a mean number of 0.054 items per gram (Barboza et al. 2020). Based on the fish muscle data and the recommended human consumption of fish per capita in selected European and American countries, researchers were able to estimate that adults potentially consume 518 – 3078 MP items/year/capita. These numbers are considerably smaller than the 39 000 mentioned earlier, however the figure was based on data from one type of fish; consumption of other products containing MPs were not considered.

Although most studies look at non-human MP ingestion, they can be used to look at the effects this might have on tissues and organs. A study on tissue accumulation of polystyrene in zebrafish found 5µm diameter MPs in the gills, liver and gut and 20µm diameter accumulation in just the gills and gut. This caused inflammation and lipid accumulation. They also found that exposure to MPs induced alterations of metabolic profiles in the liver and disturbed lipid and energy metabolism (Lu et al. 2016).

In a separate study, Zebrafish were exposed to three shapes of MPs (bead, fragment and fibre). There was accumulation of the MPs in the gut, with the fibre shape resulting in the

more severe intestinal toxicity than fragments and beads. The accumulation caused mucosal damage, increased permeability, inflammation, metabolism disruption and microbiota dysbiosis (Qiao et al. 2019). A different study found that fish with MPs had significantly higher lipid peroxidation levels in the brain, gills and dorsal muscle and increased brain acetylcholinesterase activity than fish with no MPs, suggesting lipid oxidative damage (Barboza et al. 2020). Medaka fish larvae and juveniles were fed food spiked with environmental MPs in a different study. They found that in those fed the spiked food it could cause death, decreased head/body ratios, increased Ethoxyresorufin-O-deethylase (EROD) activity, DNA breaks and alterations to swimming behaviour (Pannetier et al. 2020).

There was also a study that looked at polystyrene MPs in mice based on toxicity-based toxicokinetic/toxicodynamic modelling to quantify organ bioaccumulation and biomarker responses. The gut had the highest bioaccumulation factor and overall the smaller MP size (5µm) exhibited higher values compared to the larger MP size (20µm) (Yang et al. 2019b). Another study on mice and polystyrene MPs looked at tissue distribution, accumulation and tissue-specific health risks. They were found to accumulate in the liver, kidneys and gut, depending on particle size. Biochemical biomarkers suggested exposure induced disturbance of energy and lipid metabolism as well as oxidative stress (Deng et al. 2017). There is however, an editorial that is critical of this Deng et al. study, suggesting that the conclusion of health effects by MPs is not sufficiently supported by the data presented (Braeuning 2019). They also question the values given for the accumulation of plastic in organs as the figures seem to far exceed the doses administered.

A different study looked at the exposure to differing amounts of polyethylene MPs in mice. The high concentration of MPs increased the numbers of gut microbial species, bacterial

abundance and flora diversity. Serum levels of interleukin 1a in all feeding groups were significantly greater than in the blank group. The intestines of mice fed high concentrations of MPs showed inflammation and higher TLR4, AP-1 and IRF5 expression (Li et al. 2020). A contrasting study looked at the intestinal particle uptake and health-related effects of oral polystyrene in vitro and in vivo. It suggested that oral exposure to polystyrene in the conditions tested did not pose any relevant acute health risks to mammals; no inflammatory response or lesions and no interference of macrophages (Stock et al. 2019).

There should also be consideration for the effects of MPs on offspring during gestation. In a recent study, pregnant mice were exposed to polystyrene MPs in their drinking water and the offspring observed and tested. They found that there was no significant effect on the offspring's growth, however there was indication of the offspring having fatty acid metabolic disorders, which was related to the MP particle size (Luo et al. 2019). Another study used an ex vivo human placental perfusion model and fluorescently labelled polystyrene beads from 50nm – 500nm in diameter to see if the particles could cross the placental barrier and affect the foetus (Wick et al. 2010). They found that a diameter of up to 240nm was able to cross the placental barrier but did not affect the viability of the placental explant.

As mentioned previously there is a wide occurrence and reporting of MP ingestion by aquatic fauna but there are questions as to how much of this actually transfers to humans in terms of ingestion. Most studies that look at the health effects are produced in laboratory conditions which are less relevant when applied to the environment. In particular there is a need to be realistic with the concentrations of MPs used and aligning them with what would most likely be found in the environment (de Sá et al. 2018; Wang et al. 2019).

## Prosthetics

There has been limited studies on MPs generated by wear and corrosion of joint replacement prostheses, however one published in 2000, details identification of metallic and polyethylene particles from post-mortems in 29 patients (Urban et al. 2000). They found polyethylene particles in the para-aortic lymph nodes in 68% of 28 patients and in the liver or spleen of 14% of 29 patients. The majority of the particles were less than 1µm in size and mostly in low concentrations, with little pathological importance. However, in one case granulomas formed in the liver, spleen and abdominal lymph nodes in response to heavy accumulation of wear debris from a hip prosthesis. Additional studies related to plastic prosthesis particle contamination have been carried out in vitro or using in vivo models and seek to explore the health effects of the wear debris. An in vitro study found that when adding polymethylmethacrylate (PMMA) particles to each developmental stage of osteoclasts there was an increase in bone resorption in mature osteoclasts (H. Zhang et al. 2008). It is thought that inflammation is caused by PMMA particles increasing osteoclast formation resulting in prosthetic failure. Similarly, ultra-high molecular weight polyethylene (UHMWPE) particles were introduced to bone implants in mice and then treated with erythromycin 2 weeks after implantation (Markel et al. 2009). Results showed that exposure to UHMWPE particles induced inflammation and increased bone resorption, but with erythromycin treatment, this was reduced. A study in 2011 looked at toll-like receptors (TLRs) and their role in recognising orthopaedic implant wear-debris particles, ultimately causing inflammation (Pearl et al. 2011). They found that TLRs signal through myeloid differentiation factor 88 (MyD88) and by inhibiting MyD88 there was a decrease in PMMA particle induced production of macrophages and therefore, inflammation.



## Biological effects

### Microorganisms

It has been shown that bacteria can rapidly colonize MP surfaces in the marine environment (Harrison et al. 2014; Wagner et al. 2014), as well as form microbial biofilms (Lobelle and Cunliffe 2011). Zettler et al. studied plastic marine debris using a scanning electron microscope (SEM) and found a diverse microbial community coined a ‘Plastisphere’ (Zettler, Mincer, and Amaral-Zettler 2013). They found that the hydrophobic surface of these plastics is ideal for microbial colonization and biofilm formation, discovering that the most abundant genus was *Vibrio*, among many others. Although there is sufficient research showing that microorganisms can colonise MPs, there is little evidence of whether or not they are capable of degrading MPs in the field. Laboratory studies, however, indicate that fungi, bacteria and biofilms are capable of degrading MPs of a variety of polymer types including polyethylene, polystyrene and polylactic acid (Yuan et al. 2020).

Potentially pathogenic *Vibrio* spp. were found to be present on floating MPs in water samples from the North and Baltic Sea, which suggests MPs could function as vectors for the dispersal of pathogens (Kirstein et al. 2016). The *Vibrio* spp. Pathogen could cause serious infections in humans if ingested (Morris Jr. and Acheson 2003), illustrating that the presence of MPs in seafood is an area to be investigated further.

There is some evidence to suggest that interactions between MPs, microorganisms and gut microbiota could lead to health implications (Lu et al. 2019). It is known that gut microbiota plays an important role in the hosts health and it is also known that MPs can carry potential pesticides, fungicides and pathogens. Once ingested these hitchhikers may affect health by changing the composition of gut microbiota (Lu et al. 2019).

424

425 A further area of interest is the antibiotic resistant bacteria (ARB) that has been found on  
426 some MPs (Yang et al. 2019a). A recent study found ARB counts on MPs were 100 – 500  
427 times higher than those in water and the ratios of ARB to total bacteria from MPs were higher  
428 than those in water (Zhang et al. 2020c). They also looked at multi-antibiotic resistant  
429 bacteria (MARB) and found penicillin, sulfafurazole, erythromycin and tetracycline resistant  
430 bacteria, accounted for 25.4% on MPs compared with 23.9% in water. Further studies  
431 looking at antibiotic resistant genes (ARG) showed a detection rate up to 80% on MPs and  
432 65% in water (Zhang et al. 2020c). It was concluded that MPs can provide a beneficial  
433 surface for ARB to form a biofilm and facilitate horizontal gene transfer, which they would  
434 otherwise be unable to do in water alone and this could lead to the enrichment of superbugs.

435

## 436 **Conclusion**

437 There is a growing body of literature demonstrating that the atmosphere and human food and  
438 water sources are contaminated by MPs and may, therefore, be inhaled or ingested by  
439 humans. Studies using model organisms indicate that ingestion of microplastics might cause  
440 harm to organisms via their physical presence (abrasive effects leading to inflammation,  
441 oxidative stress and cytotoxicity), their chemical burden (leaching of additives or adsorbed  
442 chemicals from the environment causing reproductive and developmental toxicity or  
443 invoking an immune response) or their microbial communities (pathogens causing infection,  
444 gut dysbiosis or antimicrobial resistant microbes entering the body). In addition, inhalation of  
445 plastic microfibres has become a key research focus with recent estimates suggesting the  
446 general population inhales hundreds of plastic fibres each day. Correlative research links  
447 inhalation of plastic fibres to respiratory disease, inflammation and oxidative stress making  
448 inhalation of microfibres a key area of concern given the growing dominance of synthetic

449 fibres in the clothing industry. The actual concentrations of inhaled and ingested  
450 microplastics that are accumulated within the human body are, however, not yet known.  
451 There is still a dearth of data on the direct human health implications and future work should  
452 target the direct effects of MPs on human health by focusing on inflammation and cellular  
453 damage at concentrations realistically reflecting environmental exposure.  
454

## References

- Abb, M., T. Heinrich, E. Sorkau, and W. Lorenz. 2009. "Phthalates in House Dust." *Environment International* 35 (6): 965–70. <https://doi.org/10.1016/J.ENVINT.2009.04.007>.
- Abbasi, S., B. Keshavarzi, F. Moore, A. Turner, F.J. Kelly, A. Dominguez, and N. Jaafarzadeh. 2019. "Distribution and Potential Health Impacts of Microplastics and Microrubbers in Air and Street Dusts from Asaluyeh County, Iran." *Environmental Pollution* 244 (January): 153–64. <https://doi.org/10.1016/J.ENVPOL.2018.10.039>.
- Abidli, S., J.C. Antunes, J.L. Ferreira, Y. Lahbib, P. Sobral, and N. Trigui El Menif. 2018. "Microplastics in Sediments from the Littoral Zone of the North Tunisian Coast (Mediterranean Sea)." *Estuarine, Coastal and Shelf Science* 205 (May): 1–9. <https://doi.org/10.1016/J.ECSS.2018.03.006>.
- Ait Bamai, Y., A. Araki, T. Kawai, T. Tsuboi, I. Saito, E. Yoshioka, A. Kanazawa, et al. 2014. "Associations of Phthalate Concentrations in Floor Dust and Multi-Surface Dust with the Interior Materials in Japanese Dwellings." *Science of The Total Environment* 468–469: 147–57. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2013.07.107>.
- Ait Bamai, Y., E. Shibata, I. Saito, A. Araki, A. Kanazawa, K. Morimoto, K. Nakayama, et al. 2014. "Exposure to House Dust Phthalates in Relation to Asthma and Allergies in Both Children and Adults." *Science of The Total Environment* 485–486: 153–63. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2014.03.059>.
- Allen, S., D. Allen, V.R. Phoenix, G. Le Roux, P. Durántez Jiménez, A. Simonneau, S. Binet, and D. Galop. 2019. "Atmospheric Transport and Deposition of Microplastics in a Remote Mountain Catchment." *Nature Geoscience* 12 (5): 339–44. <https://doi.org/10.1038/s41561-019-0335-5>.
- Alonso, V., V. Linares, M. Bellés, M.L Albina, A. Pujol, J.L. Domingo, and D.J. Sánchez.

480 2010. “Effects of BDE-99 on Hormone Homeostasis and Biochemical Parameters in  
 481 Adult Male Rats.” *Food and Chemical Toxicology* 48 (8): 2206–11.  
 482 <https://doi.org/https://doi.org/10.1016/j.fct.2010.05.048>.

483 Ambrosini, R., R.S. Azzoni, F. Pittino, G. Diolaiuti, A. Franzetti, and M. Parolini. 2019.  
 484 “First Evidence of Microplastic Contamination in the Supraglacial Debris of an Alpine  
 485 Glacier.” *Environmental Pollution* 253 (October): 297–301.  
 486 <https://doi.org/10.1016/J.ENVPOL.2019.07.005>.

487 Amélineau, F., D. Bonnet, O. Heitz, V. Mortreux, A.M.A. Harding, N. Karnovsky, W.  
 488 Walkusz, J. Fort, and D. Grémillet. 2016. “Microplastic Pollution in the Greenland Sea:  
 489 Background Levels and Selective Contamination of Planktivorous Diving Seabirds.”  
 490 *Environmental Pollution* 219 (December): 1131–39.  
 491 <https://doi.org/10.1016/J.ENVPOL.2016.09.017>.

492 Ashton, K., L. Holmes, and A. Turner. 2010. “Association of Metals with Plastic Production  
 493 Pellets in the Marine Environment.” *Marine Pollution Bulletin* 60 (11): 2050–55.  
 494 <https://doi.org/10.1016/J.MARPOLBUL.2010.07.014>.

495 Auta, H.S., C.U. Emenike, and S.H. Fauziah. 2017. “Distribution and Importance of  
 496 Microplastics in the Marine Environment: A Review of the Sources, Fate, Effects, and  
 497 Potential Solutions.” *Environment International* 102 (May): 165–76.  
 498 <https://doi.org/10.1016/J.ENVINT.2017.02.013>.

499 Avio CG, Gorbi S, Milan M, Benedetti M, Fattorini D, d’Errico G, Pauletto M, Bargelloni L,  
 500 Regoli F (2015) Pollutants bioavailability and toxicological risk from microplastics to  
 501 marine mussels. *Environ Pollut* 198:211–222.

502 Bakir, A., S.J. Rowland, and R.C. Thompson. 2012. “Competitive Sorption of Persistent  
 503 Organic Pollutants onto Microplastics in the Marine Environment.” *Marine Pollution*  
 504 *Bulletin* 64 (12): 2782–89. <https://doi.org/10.1016/J.MARPOLBUL.2012.09.010>.

505 Bakir, A., O'Connor, I.A., Rowland, S.J., Hendriks, A.J., and Thompson, R.C. (2016).  
506 Relative importance of microplastics as a pathway for the transfer of hydrophobic  
507 organic chemicals to marine life. *Environ. Pollut.*, 219:56-65.

508 Bank, M.S., and S.V. Hansson. 2019. "The Plastic Cycle: A Novel and Holistic Paradigm for  
509 the Anthropocene." *Environmental Science & Technology* 53 (13): 7177–79.  
510 <https://doi.org/10.1021/acs.est.9b02942>.

511 Barboza, L.G.A., A. Dick Vethaak, B. Lavorante, A.K. Lundebye, and L. Guilhermino. 2018.  
512 "Marine Microplastic Debris: An Emerging Issue for Food Security, Food Safety and  
513 Human Health." *Marine Pollution Bulletin* 133 (August): 336–48.  
514 <https://doi.org/10.1016/J.MARPOLBUL.2018.05.047>.

515 Barboza, L.G.A., C. Lopes, P. Oliveira, F. Bessa, V. Otero, B. Henriques, J. Raimundo, M.  
516 Caetano, Carlos Vale, and Lúcia Guilhermino. 2020. "Microplastics in Wild Fish from  
517 North East Atlantic Ocean and Its Potential for Causing Neurotoxic Effects, Lipid  
518 Oxidative Damage, and Human Health Risks Associated with Ingestion Exposure."  
519 *Science of The Total Environment* 717: 134625.  
520 <https://doi.org/https://doi.org/10.1016/j.scitotenv.2019.134625>.

521 Bellés, M., Vi. Alonso, V. Linares, M.L. Albina, J.J. Sirvent, J.L. Domingo, and D.J.  
522 Sánchez. 2010. "Behavioral Effects and Oxidative Status in Brain Regions of Adult Rats  
523 Exposed to BDE-99." *Toxicology Letters* 194 (1): 1–7.  
524 <https://doi.org/https://doi.org/10.1016/j.toxlet.2010.01.010>.

525 Besseling, E., A. Wegner, E. Foekema, M. van den Heuvel-Greve, and A. Koelmans. 2013.  
526 "Effects of Microplastic on Fitness and PCB Bioaccumulation by the Lugworm  
527 *Arenicola Marina* (L.)." *Environmental Science & Technology* 47 (1): 593–600.  
528 <https://doi.org/10.1021/es302763x>.

529 Boucher, J., and G. Billard. 2019. "The Challenges of Measuring Plastic Pollution." *Field*

530 *Actions Science Reports*, no. Special Issue 19: 68–75.

531 <https://journals.openedition.org/factsreports/5319#tocto1n6>.

532 Boucher, J., and D. Friot. 2017. *Primary Microplastics in the Oceans: A Global Evaluation*

533 *of Sources*. IUCN Gland, Switzerland.

534 Braeuning, A. 2019. “Uptake of Microplastics and Related Health Effects: A Critical

535 Discussion of Deng et Al., Scientific Reports 7:46687, 2017.” *Archives of Toxicology* 93

536 (1): 219–20. <https://doi.org/10.1007/s00204-018-2367-9>.

537 Brown, D.M., M.R. Wilson, W. MacNee, V. Stone, and K. Donaldson. 2001. “Size-

538 Dependent Proinflammatory Effects of Ultrafine Polystyrene Particles: A Role for

539 Surface Area and Oxidative Stress in the Enhanced Activity of Ultrafines.” *Toxicology*

540 *and Applied Pharmacology* 175 (3): 191–99. <https://doi.org/10.1006/TAAP.2001.9240>.

541 Browne, M.A., P. Crump, S.J. Niven, E. Teuten, A. Tonkin, T. Galloway, and R. Thompson.

542 2011. “Accumulation of Microplastic on Shorelines Worldwide: Sources and Sinks.”

543 *Environmental Science & Technology* 45 (21): 9175–79.

544 <https://doi.org/10.1021/es201811s>.

545 Browne, M.A., S.J. Niven, T.S. Galloway, S.J. Rowland, and R.C. Thompson. 2013.

546 “Microplastic Moves Pollutants and Additives to Worms, Reducing Functions Linked to

547 Health and Biodiversity.” *Current Biology* 23 (23): 2388–92.

548 <https://doi.org/10.1016/j.cub.2013.10.012>.

549 Cai, L., J. Wang, J. Peng, Z. Tan, Z. Zhan, X. Tan, and Q. Chen. 2017. “Characteristic of

550 Microplastics in the Atmospheric Fallout from Dongguan City, China: Preliminary

551 Research and First Evidence.” *Environmental Science and Pollution Research* 24 (32):

552 24928–35. <https://doi.org/10.1007/s11356-017-0116-x>.

553 Catarino, A.I., V. Macchia, W.G. Sanderson, R.C. Thompson, and T.B. Henry. 2018. “Low

554 Levels of Microplastics (MP) in Wild Mussels Indicate That MP Ingestion by Humans

555 Is Minimal Compared to Exposure via Household Fibres Fallout during a Meal.”  
 556 *Environmental Pollution* 237 (June): 675–84.  
 557 <https://doi.org/10.1016/J.ENVPOL.2018.02.069>.

558 Choudhary, D., C. Kurien, and A.K. Srivastava. 2020. “Microplastic Contamination and Life  
 559 Cycle Assessment of Bottled Drinking Water.” In *Advances in Water Pollution*  
 560 *Monitoring and Control*, edited by Nihal Anwar Siddiqui, S M Tauseef, and Rajendra  
 561 Dobhal, 41–48. Singapore: Springer Singapore.

562 Cox, K.D., G.A. Covernton, H.L. Davies, J.F. Dower, F. Juanes, and S.E. Dudas. 2019.  
 563 “Human Consumption of Microplastics.” *Environmental Science & Technology* 53 (12):  
 564 7068–74. <https://doi.org/10.1021/acs.est.9b01517>.

565 Crawford, C.B., and B. Quinn. 2017. “The Interactions of Microplastics and Chemical  
 566 Pollutants.” *Microplastic Pollutants*, January, 131–57. [https://doi.org/10.1016/B978-0-](https://doi.org/10.1016/B978-0-12-809406-8.00006-2)  
 567 [12-809406-8.00006-2](https://doi.org/10.1016/B978-0-12-809406-8.00006-2).

568 Dehghani, S., F. Moore, and R. Akhbarizadeh. 2017. “Microplastic Pollution in Deposited  
 569 Urban Dust, Tehran Metropolis, Iran.” *Environmental Science and Pollution Research*  
 570 24 (25): 20360–71. <https://doi.org/10.1007/s11356-017-9674-1>.

571 Deng, Y., Y. Zhang, B. Lemos, and H. Ren. 2017. “Tissue Accumulation of Microplastics in  
 572 Mice and Biomarker Responses Suggest Widespread Health Risks of Exposure.”  
 573 *Scientific Reports* 7 (1): 46687. <https://doi.org/10.1038/srep46687>.

574 Domingo, J.L. 2012. “Polybrominated Diphenyl Ethers in Food and Human Dietary  
 575 Exposure: A Review of the Recent Scientific Literature.” *Food and Chemical*  
 576 *Toxicology* 50 (2): 238–49. [https://doi.org/https://doi.org/10.1016/j.fct.2011.11.004](https://doi.org/10.1016/j.fct.2011.11.004).

577 Driedger, A.G.J., H.H. Dürr, K. Mitchell, and P. Van Cappellen. 2015. “Plastic Debris in the  
 578 Laurentian Great Lakes: A Review.” *Journal of Great Lakes Research* 41 (1): 9–19.  
 579 <https://doi.org/10.1016/J.JGLR.2014.12.020>.



580 Dris, R., J. Gasperi, C. Mirande, C. Mandin, M. Guerrouache, V. Langlois, and B. Tassin.  
581 2017. “A First Overview of Textile Fibers, Including Microplastics, in Indoor and  
582 Outdoor Environments.” *Environmental Pollution* 221 (February): 453–58.  
583 <https://doi.org/10.1016/J.ENVPOL.2016.12.013>.

584 Dris, R., J. Gasperi, V. Rocher, M. Saad, N. Renault, and B. Tassin. 2015. “Microplastic  
585 Contamination in an Urban Area: A Case Study in Greater Paris.” *Environmental*  
586 *Chemistry* 12 (5): 592–99. <https://doi.org/10.1071/EN14167>.

587 Eyles, J.E., V.W. Bramwell, E.D. Williamson, and H.O. Alpar. 2001. “Microsphere  
588 Translocation and Immunopotential in Systemic Tissues Following Intranasal  
589 Administration.” *Vaccine* 19 (32): 4732–42. [https://doi.org/10.1016/S0264-](https://doi.org/10.1016/S0264-410X(01)00220-1)  
590 [410X\(01\)00220-1](https://doi.org/10.1016/S0264-410X(01)00220-1).

591 Fang, C., R. Zheng, H. Chen, F. Hong, L. Lin, H. Lin, H. Guo, et al. 2019. “Comparison of  
592 Microplastic Contamination in Fish and Bivalves from Two Major Cities in Fujian  
593 Province, China and the Implications for Human Health.” *Aquaculture* 512 (October):  
594 734322. <https://doi.org/10.1016/J.AQUACULTURE.2019.734322>.

595 Franeker, J.A. van, C. Blaize, J. Danielsen, K. Fairclough, J. Gollan, N. Guse, P.L. Hansen, et  
596 al. 2011. “Monitoring Plastic Ingestion by the Northern Fulmar *Fulmarus Glacialis* in  
597 the North Sea.” *Environmental Pollution* 159 (10): 2609–15.  
598 <https://doi.org/10.1016/J.ENVPOL.2011.06.008>.

599 Frias, J.P.G.L., P. Sobral, and A.M. Ferreira. 2010. “Organic Pollutants in Microplastics from  
600 Two Beaches of the Portuguese Coast.” *Marine Pollution Bulletin* 60 (11): 1988–92.  
601 <https://doi.org/10.1016/J.MARPOLBUL.2010.07.030>.

602 Gallagher, L.G., W. Li, R.M. Ray, M.E. Romano, K.J. Wernli, D. Gao, D.B. Thomas, and H.  
603 Checkoway. 2015. “Occupational Exposures and Risk of Stomach and Esophageal  
604 Cancers: Update of a Cohort of Female Textile Workers in Shanghai, China.” *American*

605 *Journal of Industrial Medicine* 58 (3): 267–75. <https://doi.org/10.1002/ajim.22412>.

606 Galloway, T.S. 2015. “Micro- and Nano-Plastics and Human Health BT - Marine

607 Anthropogenic Litter.” In , edited by Melanie Bergmann, Lars Gutow, and Michael

608 Klages, 343–66. Cham: Springer International Publishing. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-3-319-16510-3_13)

609 [3-319-16510-3\\_13](https://doi.org/10.1007/978-3-319-16510-3_13).

610 Gasperi, J., S.L. Wright, R. Dris, F. Collard, C. Mandin, M. Guerrouache, V. Langlois, F.J.

611 Kelly, and B. Tassin. 2018. “Microplastics in Air: Are We Breathing It In?” *Current*

612 *Opinion in Environmental Science & Health* 1 (February): 1–5.

613 <https://doi.org/10.1016/J.COESH.2017.10.002>.

614 Geyer, R., J.R. Jambeck, and K.L. Law. 2017. “Production, Use, and Fate of All Plastics Ever

615 Made.” *Science Advances* 3 (7): e1700782. <https://doi.org/10.1126/sciadv.1700782>.

616 Gouin, T., N. Roche, R. Lohmann, and G. Hodges. 2011. “A Thermodynamic Approach for

617 Assessing the Environmental Exposure of Chemicals Absorbed to Microplastic.”

618 *Environmental Science & Technology* 45 (4): 1466–72.

619 <https://doi.org/10.1021/es1032025>.

620 Harrison, J.P., M. Schratzberger, M. Sapp, and A.M. Osborn. 2014. “Rapid Bacterial

621 Colonization of Low-Density Polyethylene Microplastics in Coastal Sediment

622 Microcosms.” *BMC Microbiology* 14 (1): 232. [https://doi.org/10.1186/s12866-014-](https://doi.org/10.1186/s12866-014-0232-4)

623 [0232-4](https://doi.org/10.1186/s12866-014-0232-4).

624 Hartmann, N.B., T. Hüffer, R.C. Thompson, M. Hassellöv, A. Verschoor, A.E. Dagaard, S.

625 Rist, et al. 2019. “Are We Speaking the Same Language? Recommendations for a

626 Definition and Categorization Framework for Plastic Debris.” *Environmental Science &*

627 *Technology* 53 (3): 1039–47. <https://doi.org/10.1021/acs.est.8b05297>.

628 Högberg, J., A. Hanberg, M. Berglund, S. Skerfving, M. Remberger, A.M Calafat, A. Falk

629 Filipsson, et al. 2008. “Phthalate Diesters and Their Metabolites in Human Breast Milk,

630 Blood or Serum, and Urine as Biomarkers of Exposure in Vulnerable Populations.”

631 *Environmental Health Perspectives* 116 (3): 334–39. <https://doi.org/10.1289/ehp.10788>.

632 Holmes, L.A., and R.C. Thompson. 2014. “Interactions between Trace Metals and Plastic

633 Production Pellets under Estuarine Conditions.” *Marine Chemistry* 167 (December): 25–

634 32. <https://doi.org/10.1016/J.MARCHEM.2014.06.001>.

635 Holmes, L.A., A. Turner, and R.C. Thompson. 2012. “Adsorption of Trace Metals to Plastic

636 Resin Pellets in the Marine Environment.” *Environmental Pollution* 160 (January): 42–

637 48. <https://doi.org/10.1016/J.ENVPOL.2011.08.052>.

638 Horton, A.A., and S.J. Dixon. 2018. “Microplastics: An Introduction to Environmental

639 Transport Processes.” *WIREs Water* 5 (2): e1268. <https://doi.org/10.1002/wat2.1268>.

640 Horton, A.A., C. Svendsen, R. Williams, D.J. Spurgeon, and E. Lahive. 2017. “Large

641 Microplastic Particles in Sediments of Tributaries of the River Thames, UK –

642 Abundance, Sources and Methods for Effective Quantification.” *Marine Pollution*

643 *Bulletin* 114 (1): 218–26. <https://doi.org/10.1016/J.MARPOLBUL.2016.09.004>.

644 Huang, Y., Q. Liu, W. Jia, C. Yan, and J. Wang. 2020. “Agricultural Plastic Mulching as a

645 Source of Microplastics in the Terrestrial Environment.” *Environmental Pollution* 260:

646 114096. <https://doi.org/https://doi.org/10.1016/j.envpol.2020.114096>.

647 Huang, Y., X. Qing, W. Wang, G. Han, and J. Wang. 2020. “Mini-Review on Current Studies

648 of Airborne Microplastics: Analytical Methods, Occurrence, Sources, Fate and Potential

649 Risk to Human Beings.” *TrAC Trends in Analytical Chemistry* 125 (April): 115821.

650 <https://doi.org/10.1016/J.TRAC.2020.115821>.

651 Huerta Lwanga, E., H. Gertsen, H. Gooren, P. Peters, T. Salánki, M. van der Ploeg, E.

652 Besseling, A.A. Koelmans, and Violette Geissen. 2016. “Microplastics in the Terrestrial

653 Ecosystem: Implications for Lumbricus Terrestris (Oligochaeta, Lumbricidae).”

654 *Environmental Science & Technology* 50 (5): 2685–91.

655 <https://doi.org/10.1021/acs.est.5b05478>.

656 Hwang, J., D. Choi, S. Han, J. Choi, and J. Hong. 2019. “An Assessment of the Toxicity of  
657 Polypropylene Microplastics in Human Derived Cells.” *Science of The Total*  
658 *Environment* 684 (September): 657–69.  
659 <https://doi.org/10.1016/J.SCITOTENV.2019.05.071>.

660 Jambeck, J.R., R. Geyer, C. Wilcox, T. Siegler, M. Perryman, A. Andrady, R. Narayan, and  
661 K.L. Law. 2015. “Plastic Waste Inputs from Land into the Ocean.” *Science* 347 (6223):  
662 768 LP – 771. <https://doi.org/10.1126/science.1260352>.

663 Jiang, C., L. Yin, Z. Li, X. Wen, X. Luo, S. Hu, H. Yang, et al. 2019. “Microplastic Pollution  
664 in the Rivers of the Tibet Plateau.” *Environmental Pollution* 249 (June): 91–98.  
665 <https://doi.org/10.1016/J.ENVPOL.2019.03.022>.

666 Jornet-Martínez, N., C. Antón-Soriano, and P. Campíns-Falcó. 2015. “Estimation of the  
667 Presence of Unmetabolized Dialkyl Phthalates in Untreated Human Urine by an On-  
668 Line Miniaturized Reliable Method.” *Science of The Total Environment* 532  
669 (November): 239–44. <https://doi.org/10.1016/J.SCITOTENV.2015.05.124>.

670 Katyal, D., E. Kong, and J. Villanueva. 2020. “Microplastics in the Environment: Impact on  
671 Human Health and Future Mitigation Strategies.” *Environmental Health Review* 63 (1):  
672 27–31. <https://doi.org/10.5864/d2020-005>.

673 Kirstein, I., S. Kirmizi, A. Wichels, A. Garin-Fernandez, R. Erler, M. Löder, and G. Gerdtz.  
674 2016. “Dangerous Hitchhikers? Evidence for Potentially Pathogenic *Vibrio* Spp. on  
675 Microplastic Particles.” *Marine Environmental Research* 120 (September): 1–8.  
676 <https://doi.org/10.1016/j.marenvres.2016.07.004>.

677 Koelmans, A.A., E. Besseling, A. Wegner, and E.M. Foekema. 2013. “Plastic as a Carrier of  
678 POPs to Aquatic Organisms: A Model Analysis.” *Environmental Science & Technology*  
679 47 (14): 7812–20. <https://doi.org/10.1021/es401169n>.

680 Koelmans, A.A., Bakir, A., Burton, G.A., and Janssen C.R. (2016). Microplastic as a vector  
 681 for chemicals in the aquatic environment. Critical review and model-supported  
 682 reinterpretation of empirical studies. *Environ. Sci. Technol.* 50, 3315-3326.  
 683 Kosuth, M., S.A. Mason, and E.V. Wattenberg. 2018. "Anthropogenic Contamination of Tap  
 684 Water, Beer, and Sea Salt." *PLOS ONE* 13 (4): e0194970.  
 685 <https://doi.org/10.1371/journal.pone.0194970>.  
 686 Latini, G., C. De Felice, G. Presta, A. Del Vecchio, I. Paris, F. Ruggieri, and P. Mazzeo.  
 687 2003. "In Utero Exposure to Di-(2-Ethylhexyl)Phthalate and Duration of Human  
 688 Pregnancy." *Environmental Health Perspectives* 111 (14): 1783–85.  
 689 <https://doi.org/10.1289/ehp.6202>.  
 690 Lavers, J.L., A.L. Bond, and I. Hutton. 2014. "Plastic Ingestion by Flesh-Footed Shearwaters  
 691 (Puffinus Carneipes): Implications for Fledgling Body Condition and the Accumulation  
 692 of Plastic-Derived Chemicals." *Environmental Pollution* 187 (April): 124–29.  
 693 <https://doi.org/10.1016/J.ENVPOL.2013.12.020>.  
 694 Law, K.L., and R.C. Thompson. 2014. "Microplastics in the Seas." *Science* 345 (6193): 144  
 695 LP – 145. <https://doi.org/10.1126/science.1254065>.  
 696 Lehner, R., C. Weder, A. Petri-Fink, and B. Rothen-Rutishauser. 2019. "Emergence of  
 697 Nanoplastic in the Environment and Possible Impact on Human Health." *Environmental*  
 698 *Science & Technology* 53 (4): 1748–65. <https://doi.org/10.1021/acs.est.8b05512>.  
 699 Li, B., Y. Ding, X. Cheng, D. Sheng, Z. Xu, Q. Rong, Y. Wu, H. Zhao, Xiaofei Ji, and Ying  
 700 Zhang. 2020. "Polyethylene Microplastics Affect the Distribution of Gut Microbiota and  
 701 Inflammation Development in Mice." *Chemosphere* 244 (April): 125492.  
 702 <https://doi.org/10.1016/J.CHEMOSPHERE.2019.125492>.  
 703 Li, J., H. Liu, and J.P. Chen. 2018. "Microplastics in Freshwater Systems: A Review on  
 704 Occurrence, Environmental Effects, and Methods for Microplastics Detection." *Water*

705        *Research* 137 (June): 362–74. <https://doi.org/10.1016/J.WATRES.2017.12.056>.

706    Li, J., X. Qu, L. Su, W. Zhang, D. Yang, P. Kolandhasamy, D. Li, and H. Shi. 2016.

707        “Microplastics in Mussels along the Coastal Waters of China.” *Environmental Pollution*

708        214 (July): 177–84. <https://doi.org/10.1016/J.ENVPOL.2016.04.012>.

709    Li, Jia, Kaina Zhang, and Hua Zhang. 2018. “Adsorption of Antibiotics on Microplastics.”

710        *Environmental Pollution* 237: 460–67.

711        <https://doi.org/https://doi.org/10.1016/j.envpol.2018.02.050>.

712    Linares, V., M. Bellés, and J.L. Domingo. 2015. “Human Exposure to PBDE and Critical

713        Evaluation of Health Hazards.” *Archives of Toxicology* 89 (3): 335–56.

714        <https://doi.org/10.1007/s00204-015-1457-1>.

715    Lithner, D., J. Damberg, G. Dave, and Å. Larsson. 2009. “Leachates from Plastic Consumer

716        Products - Screening for Toxicity with *Daphnia Magna*.” *Chemosphere*.

717        <https://doi.org/10.1016/j.chemosphere.2008.11.022>.

718    Lithner, D., Å. Larsson, and G. Dave. 2011. “Environmental and Health Hazard Ranking and

719        Assessment of Plastic Polymers Based on Chemical Composition.” *Science of The Total*

720        *Environment* 409 (18): 3309–24. <https://doi.org/10.1016/J.SCITOTENV.2011.04.038>.

721    Liu, K., X. Wang, T. Fang, P. Xu, L. Zhu, and D. Li. 2019. “Source and Potential Risk

722        Assessment of Suspended Atmospheric Microplastics in Shanghai.” *Science of The*

723        *Total Environment* 675 (July): 462–71.

724        <https://doi.org/10.1016/J.SCITOTENV.2019.04.110>.

725    Liu, K., X. Wang, N. Wei, Z. Song, and D. Li. 2019. “Accurate Quantification and Transport

726        Estimation of Suspended Atmospheric Microplastics in Megacities: Implications for

727        Human Health.” *Environment International* 132: 105127.

728        <https://doi.org/https://doi.org/10.1016/j.envint.2019.105127>.

729    Liu, K., T. Wu, X. Wang, Z. Song, C. Zong, N. Wei, and D. Li. 2019. “Consistent Transport

730 of Terrestrial Microplastics to the Ocean through Atmosphere.” *Environmental Science*  
731 & *Technology* 53 (18): 10612–19. <https://doi.org/10.1021/acs.est.9b03427>.

732 Lobelle, D., and M. Cunliffe. 2011. “Early Microbial Biofilm Formation on Marine Plastic  
733 Debris.” *Marine Pollution Bulletin* 62 (1): 197–200.  
734 <https://doi.org/10.1016/J.MARPOLBUL.2010.10.013>.

735 Lu, J., Wu, J., Wu, J., Zhang, C., and Luo, Y. 2020. “Adsorption and Desorption of Steroid  
736 Hormones by Microplastics in Seawater.” *Bulletin of Environmental Contamination and*  
737 *Toxicology*. <https://doi.org/10.1007/s00128-020-02784-2>.

738 Lu, L., T. Luo, Y. Zhao, C. Cai, Z. Fu, and Y. Jin. 2019. “Interaction between Microplastics  
739 and Microorganism as Well as Gut Microbiota: A Consideration on Environmental  
740 Animal and Human Health.” *Science of The Total Environment* 667 (June): 94–100.  
741 <https://doi.org/10.1016/J.SCITOTENV.2019.02.380>.

742 Lu, Y., Y. Zhang, Y. Deng, W. Jiang, Y. Zhao, J. Geng, L. Ding, and H. Ren. 2016. “Uptake  
743 and Accumulation of Polystyrene Microplastics in Zebrafish (*Danio Rerio*) and Toxic  
744 Effects in Liver.” *Environmental Science & Technology* 50 (7): 4054–60.  
745 <https://doi.org/10.1021/acs.est.6b00183>.

746 Luo, T., Y. Zhang, C. Wang, X. Wang, J. Zhou, M. Shen, Y. Zhao, Z. Fu, and Yuanxiang Jin.  
747 2019. “Maternal Exposure to Different Sizes of Polystyrene Microplastics during  
748 Gestation Causes Metabolic Disorders in Their Offspring.” *Environmental Pollution* 255  
749 (December): 113122. <https://doi.org/10.1016/J.ENVPOL.2019.113122>.

750 Lusher, A.L., M. McHugh, and R.C. Thompson. 2013. “Occurrence of Microplastics in the  
751 Gastrointestinal Tract of Pelagic and Demersal Fish from the English Channel.” *Marine*  
752 *Pollution Bulletin* 67 (1–2): 94–99.  
753 <https://doi.org/10.1016/J.MARPOLBUL.2012.11.028>.

754 Main, K., G. Mortensen, M. Kaleva, K. Boisen, I. Damgaard, M. Chellakooty, I. Schmidt, et

755 al. 2006. “Human Breast Milk Contamination with Phthalates and Alterations of  
756 Endogenous Reproductive Hormones in Infants Three Months of Age.” *Environmental*  
757 *Health Perspectives* 114 (2): 270–76. <https://doi.org/10.1289/ehp.8075>.

758 Markel, D., R. Zhang, T. Shi, M. Hawkins, and W. Ren. 2009. “Inhibitory Effects of  
759 Erythromycin on Wear Debris-Induced VEGF/Flt-1 Gene Production and Osteolysis.”  
760 *Inflammation Research* 58 (7): 413. <https://doi.org/10.1007/s00011-009-0007-9>.

761 Morris Jr., J.G., and D. Acheson. 2003. “Cholera and Other Types of Vibriosis: A Story of  
762 Human Pandemics and Oysters on the Half Shell.” *Clinical Infectious Diseases* 37 (2):  
763 272–80. <https://doi.org/10.1086/375600>.

764 Oliveri Conti, G., M. Ferrante, M. Banni, C. Favara, I. Nicolosi, A. Cristaldi, M. Fiore, and P.  
765 Zuccarello. 2020. “Micro- and Nano-Plastics in Edible Fruit and Vegetables. The First  
766 Diet Risks Assessment for the General Population.” *Environmental Research* 187:  
767 109677. <https://doi.org/https://doi.org/10.1016/j.envres.2020.109677>.

768 Pannetier, P., B. Morin, F. Le Bihanic, L. Dubreil, C. Clérandeau, F. Chouvellon, K. Van  
769 Arkel, M. Danion, and J. Cachot. 2020. “Environmental Samples of Microplastics  
770 Induce Significant Toxic Effects in Fish Larvae.” *Environment International* 134  
771 (January): 105047. <https://doi.org/10.1016/J.ENVINT.2019.105047>.

772 Pauly, J.L., S.J. Stegmeier, H.A. Allaart, R.T. Cheney, P.J. Zhang, A.G. Mayer, and R.J.  
773 Streck. 1998. “Inhaled Cellulosic and Plastic Fibers Found in Human Lung Tissue.”  
774 *Cancer Epidemiology Biomarkers & Prevention* 7 (5): 419 LP – 428.  
775 <http://cebp.aacrjournals.org/content/7/5/419.abstract>.

776 Pearl, J., T. Ma, A.R. Irani, Z. Huang, W.H. Robinson, R.L. Smith, and S.B. Goodman. 2011.  
777 “Role of the Toll-like Receptor Pathway in the Recognition of Orthopedic Implant  
778 Wear-Debris Particles.” *Biomaterials* 32 (24): 5535–42.  
779 <https://doi.org/https://doi.org/10.1016/j.biomaterials.2011.04.046>.



780 Peeken, I., S. Primpke, B. Beyer, J. Gütermann, C. Katlein, T. Krumpen, M. Bergmann, L.  
 781 Hehemann, and Gunnar Gerds. 2018. "Arctic Sea Ice Is an Important Temporal Sink  
 782 and Means of Transport for Microplastic." *Nature Communications* 9 (1): 1505.  
 783 <https://doi.org/10.1038/s41467-018-03825-5>.  
 784 Peixoto, D., C. Pinheiro, J. Amorim, L. Oliva-Teles, L. Guilhermino, and M. Vieira. 2019.  
 785 "Microplastic Pollution in Commercial Salt for Human Consumption: A Review."  
 786 *Estuarine, Coastal and Shelf Science* 219 (April): 161–68.  
 787 <https://doi.org/10.1016/J.ECSS.2019.02.018>.  
 788 Peng, X., M. Chen, S. Chen, S. Dasgupta, H. Xu, K. Ta, M. Du, J. Li, Z Guo, and S Bai.  
 789 2018. "Microplastics Contaminate the Deepest Part of the World's Ocean."  
 790 *Geochemical Perspectives Letters* 9: 1–5.  
 791 <https://doi.org/http://dx.doi.org/10.7185/geochemlet.1829>.  
 792 Peretz, J., L. Vrooman, W.A. Rieke, P.A. Hunt, S. Ehrlich, R. Hauser, V. Padmanabhan, et al.  
 793 2014. "Bisphenol A and Reproductive Health: Update of Experimental and Human  
 794 Evidence, 2007–2013." *Environmental Health Perspectives* 122 (8): 775–86.  
 795 <https://doi.org/10.1289/ehp.1307728>.  
 796 Phillips, M.B., and T.H. Bonner. 2015. "Occurrence and Amount of Microplastic Ingested by  
 797 Fishes in Watersheds of the Gulf of Mexico." *Marine Pollution Bulletin* 100 (1): 264–  
 798 69. <https://doi.org/10.1016/J.MARPOLBUL.2015.08.041>.  
 799 Possatto, F.E., M. Barletta, M.F. Costa, J.A. Ivar do Sul, and D.V. Dantas. 2011. "Plastic  
 800 Debris Ingestion by Marine Catfish: An Unexpected Fisheries Impact." *Marine*  
 801 *Pollution Bulletin* 62 (5): 1098–1102.  
 802 <https://doi.org/10.1016/J.MARPOLBUL.2011.01.036>.  
 803 Prata, J.C. 2018. "Airborne Microplastics: Consequences to Human Health?" *Environmental*  
 804 *Pollution* 234 (March): 115–26. <https://doi.org/10.1016/J.ENVPOL.2017.11.043>.

805 Prata, J.C., J.P. da Costa, I. Lopes, A.C. Duarte, and T. Rocha-Santos. 2020. “Environmental  
806 Exposure to Microplastics: An Overview on Possible Human Health Effects.” *Science of*  
807 *The Total Environment* 702 (February): 134455.  
808 <https://doi.org/10.1016/J.SCITOTENV.2019.134455>.

809 Qiao, R., Y. Deng, S. Zhang, M. Wolosker, Q. Zhu, H. Ren, and Y. Zhang. 2019.  
810 “Accumulation of Different Shapes of Microplastics Initiates Intestinal Injury and Gut  
811 Microbiota Dysbiosis in the Gut of Zebrafish.” *Chemosphere* 236 (December): 124334.  
812 <https://doi.org/10.1016/J.CHEMOSPHERE.2019.07.065>.

813 Reed, S., M. Clark, R. Thompson, and K.A. Hughes. 2018. “Microplastics in Marine  
814 Sediments near Rothera Research Station, Antarctica.” *Marine Pollution Bulletin* 133  
815 (August): 460–63. <https://doi.org/10.1016/J.MARPOLBUL.2018.05.068>.

816 Reverte, I., J.L. Domingo, and M.T. Colomina. 2014. “Neurodevelopmental Effects of  
817 Decabromodiphenyl Ether (BDE-209) in APOE Transgenic Mice.” *Neurotoxicology and*  
818 *Teratology* 46: 10–17. <https://doi.org/https://doi.org/10.1016/j.ntt.2014.08.003>.

819 Ribeiro, F., J. O’Brien, T. Galloway, and K. Thomas. 2019. “Accumulation and Fate of  
820 Nano- and Micro-Plastics and Associated Contaminants in Organisms.” *TrAC Trends in*  
821 *Analytical Chemistry* 111 (February): 139–47.  
822 <https://doi.org/10.1016/J.TRAC.2018.12.010>.

823 Rios, L.M., P.R. Jones, C. Moore, and U.V. Narayan. 2010. “Quantitation of Persistent  
824 Organic Pollutants Adsorbed on Plastic Debris from the Northern Pacific Gyre’s  
825 ‘Eastern Garbage Patch.’” *Journal of Environmental Monitoring* 12 (12): 2226–36.  
826 <https://doi.org/10.1039/C0EM00239A>.

827 Rochman, C.M., B.T. Hentschel, and S.J. Teh. 2014. “Long-Term Sorption of Metals Is  
828 Similar among Plastic Types: Implications for Plastic Debris in Aquatic Environments.”  
829 *PloS One* 9 (1): e85433–e85433. <https://doi.org/10.1371/journal.pone.0085433>.

830 Rodrigues, J., A.C. Duarte, J. Santos-Echeandía, and T. Rocha-Santos. 2019. "Significance of  
831 Interactions between Microplastics and POPs in the Marine Environment: A Critical  
832 Overview." *TrAC Trends in Analytical Chemistry* 111: 252–60.  
833 <https://doi.org/https://doi.org/10.1016/j.trac.2018.11.038>.

834 Romeo, T., B. Pietro, C. Pedà, P. Consoli, F. Andaloro, and M.C. Fossi. 2015. "First  
835 Evidence of Presence of Plastic Debris in Stomach of Large Pelagic Fish in the  
836 Mediterranean Sea." *Marine Pollution Bulletin* 95 (1): 358–61.  
837 <https://doi.org/10.1016/J.MARPOLBUL.2015.04.048>.

838 Sá, L.C. de, M. Oliveira, F. Ribeiro, T. Rocha, and M.N. Futter. 2018. "Studies of the Effects  
839 of Microplastics on Aquatic Organisms: What Do We Know and Where Should We  
840 Focus Our Efforts in the Future?" *Science of The Total Environment* 645 (December):  
841 1029–39. <https://doi.org/10.1016/J.SCITOTENV.2018.07.207>.

842 Schirinzi, G.F., I. Pérez-Pomeda, J. Sanchís, C. Rossini, M. Farré, and D. Barceló. 2017.  
843 "Cytotoxic Effects of Commonly Used Nanomaterials and Microplastics on Cerebral  
844 and Epithelial Human Cells." *Environmental Research* 159 (November): 579–87.  
845 <https://doi.org/10.1016/J.ENVRES.2017.08.043>.

846 Schwabl, P., S. Köppel, P. Königshofer, T. Bucsics, M. Trauner, T. Reiberger, and B.  
847 Liebmann. 2019. "Detection of Various Microplastics in Human Stool: A Prospective  
848 Case Series." *Annals of Internal Medicine* 171 (7): 453–57.  
849 <https://doi.org/10.7326/M19-0618>.

850 Scopetani, C., A. Cincinelli, T. Martellini, E. Lombardini, A. Ciofini, A. Fortunati, V.  
851 Pasquali, S. Ciattini, and Alberto Ugolini. 2018. "Ingested Microplastic as a Two-Way  
852 Transporter for PBDEs in *Talitrus Saltator*." *Environmental Research* 167: 411–17.  
853 <https://doi.org/https://doi.org/10.1016/j.envres.2018.07.030>.

854 Shruti, V.C., F. Pérez-Guevara, I. Elizalde-Martínez, and G. Kutralam-Muniasamy. 2020.

855 “First Study of Its Kind on the Microplastic Contamination of Soft Drinks, Cold Tea and  
856 Energy Drinks - Future Research and Environmental Considerations.” *Science of The*  
857 *Total Environment* 726: 138580.  
858 <https://doi.org/https://doi.org/10.1016/j.scitotenv.2020.138580>.

859 Stock, V., L. Böhmert, E. Lisicki, R. Block, J. Cara-Carmona, L.K. Pack, R. Selb, et al. 2019.  
860 “Uptake and Effects of Orally Ingested Polystyrene Microplastic Particles in Vitro and  
861 in Vivo.” *Archives of Toxicology* 93 (7): 1817–33. [https://doi.org/10.1007/s00204-019-](https://doi.org/10.1007/s00204-019-02478-7)  
862 [02478-7](https://doi.org/10.1007/s00204-019-02478-7).

863 Talsness, C.E., A.J.M. Andrade, S.N. Kuriyama, J.A. Taylor, and F.S. vom Saal. 2009.  
864 “Components of Plastic: Experimental Studies in Animals and Relevance for Human  
865 Health.” *Philosophical Transactions of the Royal Society B: Biological Sciences* 364  
866 (1526): 2079–96. <https://doi.org/10.1098/rstb.2008.0281>.

867 Tanaka, K., H. Takada, R. Yamashita, K. Mizukawa, M. Fukuwaka, and Y. Watanuki. 2013.  
868 “Accumulation of Plastic-Derived Chemicals in Tissues of Seabirds Ingesting Marine  
869 Plastics.” *Marine Pollution Bulletin* 69 (1–2): 219–22.  
870 <https://doi.org/10.1016/J.MARPOLBUL.2012.12.010>.

871 Tang, Y., J. Rong, X. Guan, S. Zha, W. Shi, Y. Han, X. Du, F. Wu, Wei Huang, and  
872 Guangxu Liu. 2020. “Immunotoxicity of Microplastics and Two Persistent Organic  
873 Pollutants Alone or in Combination to a Bivalve Species.” *Environmental Pollution* 258:  
874 113845. <https://doi.org/https://doi.org/10.1016/j.envpol.2019.113845>.

875 Tienhoven, E.A.E. Van, D. Korbee, L. Schipper, H.W. Verharen, and W.H. De Jong. 2006.  
876 “In Vitro and in Vivo (Cyto)Toxicity Assays Using PVC and LDPE as Model  
877 Materials.” *Journal of Biomedical Materials Research Part A* 78A (1): 175–82.  
878 <https://doi.org/10.1002/jbm.a.30679>.

879 Urban, R.M., J.J. Jacobs, M.J. Tomlinson, J. Gavrilovic, J. Black, and M. Peoc’h. 2000.

880 “Dissemination of Wear Particles to the Liver, Spleen, and Abdominal Lymph Nodes of  
881 Patients with Hip or Knee Replacement\*.” *The Journal of Bone and Joint Surgery* 82  
882 (4): 457.

883 Valavanidis, A., T. Vlachogianni, K. Fiotakis, and S. Loridas. 2013. “Pulmonary Oxidative  
884 Stress, Inflammation and Cancer: Respirable Particulate Matter, Fibrous Dusts and  
885 Ozone as Major Causes of Lung Carcinogenesis through Reactive Oxygen Species  
886 Mechanisms.” *International Journal of Environmental Research and Public Health*.  
887 <https://doi.org/10.3390/ijerph10093886>.

888 Vaughan, R., S.D. Turner, and N.L. Rose. 2017. “Microplastics in the Sediments of a UK  
889 Urban Lake.” *Environmental Pollution* 229 (October): 10–18.  
890 <https://doi.org/10.1016/J.ENVPOL.2017.05.057>.

891 Velzeboer, I., C.J.A.F. Kwadijk, and A.A. Koelmans. 2014. “Strong Sorption of PCBs to  
892 Nanoplastics, Microplastics, Carbon Nanotubes, and Fullerenes.” *Environmental  
893 Science & Technology* 48 (9): 4869–76. <https://doi.org/10.1021/es405721v>.

894 Wagner, M., C. Scherer, D. Alvarez-Muñoz, N. Brennholt, X. Bourrain, S. Buchinger, E.  
895 Fries, et al. 2014. “Microplastics in Freshwater Ecosystems: What We Know and What  
896 We Need to Know.” *Environmental Sciences Europe* 26 (1): 12.  
897 <https://doi.org/10.1186/s12302-014-0012-7>.

898 Wang, T., C. Yu, Q. Chu, F. Wang, T. Lan, and J. Wang. 2020. “Adsorption Behavior and  
899 Mechanism of Five Pesticides on Microplastics from Agricultural Polyethylene Films.”  
900 *Chemosphere* 244 (April): 125491.  
901 <https://doi.org/10.1016/J.CHEMOSPHERE.2019.125491>.

902 Wang, W., H. Gao, S. Jin, R. Li, and G. Na. 2019. “The Ecotoxicological Effects of  
903 Microplastics on Aquatic Food Web, from Primary Producer to Human: A Review.”  
904 *Ecotoxicology and Environmental Safety* 173 (May): 110–17.

905 <https://doi.org/10.1016/J.ECOENV.2019.01.113>.

906 Watteau, F., M. Dignac, A. Bouchard, A. Revallier, and S. Houot. 2018. "Microplastic  
907 Detection in Soil Amended With Municipal Solid Waste Composts as Revealed by  
908 Transmission Electronic Microscopy and Pyrolysis/GC/MS." *Frontiers in Sustainable  
909 Food Systems*. <https://www.frontiersin.org/article/10.3389/fsufs.2018.00081>.

910 Wick, P., A. Malek, P. Manser, D. Meili, X. Maeder-Althaus, L. Diener, P.A. Diener, A.  
911 Zisch, Krug Harald F., and von Mandach Ursula. 2010. "Barrier Capacity of Human  
912 Placenta for Nanosized Materials." *Environmental Health Perspectives* 118 (3): 432–36.  
913 <https://doi.org/10.1289/ehp.0901200>.

914 Wright, S.L., and F.J. Kelly. 2017. "Plastic and Human Health: A Micro Issue?"  
915 *Environmental Science & Technology* 51 (12): 6634–47.  
916 <https://doi.org/10.1021/acs.est.7b00423>.

917 Wright, S.L., J. Ulke, A. Font, K.L.A. Chan, and F.J. Kelly. 2020. "Atmospheric Microplastic  
918 Deposition in an Urban Environment and an Evaluation of Transport." *Environment  
919 International* 136 (March): 105411. <https://doi.org/10.1016/J.ENVINT.2019.105411>.

920 Xiong, X., K. Zhang, X. Chen, H. Shi, Z. Luo, and C. Wu. 2018. "Sources and Distribution  
921 of Microplastics in China's Largest Inland Lake – Qinghai Lake." *Environmental  
922 Pollution* 235 (April): 899–906. <https://doi.org/10.1016/J.ENVPOL.2017.12.081>.

923 Yang, Y., G. Liu, W. Song, C. Ye, H. Lin, Z. Li, and W. Liu. 2019. "Plastics in the Marine  
924 Environment Are Reservoirs for Antibiotic and Metal Resistance Genes." *Environment  
925 International* 123: 79–86. <https://doi.org/https://doi.org/10.1016/j.envint.2018.11.061>.

926 Yang, Y.F., C.Y. Chen, T.H. Lu, and C.M. Liao. 2019. "Toxicity-Based  
927 Toxicokinetic/Toxicodynamic Assessment for Bioaccumulation of Polystyrene  
928 Microplastics in Mice." *Journal of Hazardous Materials* 366 (March): 703–13.  
929 <https://doi.org/10.1016/J.JHAZMAT.2018.12.048>.

930 Yuan, J., Ma, Y., Sun, Y., Zhou, T., Zhao, Y., Yu, F. 2020. Microbial degradation and other  
 931 environmental aspects of microplastics/plastics. *Science of The Total Environment*, 715:  
 932 136968, <https://doi.org/10.1016/j.scitotenv.2020.136968>.

933 Zarfl, C., and M. Matthies. 2010. “Are Marine Plastic Particles Transport Vectors for Organic  
 934 Pollutants to the Arctic?” *Marine Pollution Bulletin* 60 (10): 1810–14.  
 935 <https://doi.org/10.1016/J.MARPOLBUL.2010.05.026>.

936 Zettler, E.R., T.J. Mincer, and L.A. Amaral-Zettler. 2013. “Life in the ‘Plastisphere’:  
 937 Microbial Communities on Plastic Marine Debris.” *Environmental Science &*  
 938 *Technology* 47 (13): 7137–46. <https://doi.org/10.1021/es401288x>.

939 Zhang, H., B.F. Ricciardi, X. Yang, Y. Shi, N.P. Camacho, and M.P.G. Bostrom. 2008.  
 940 “Polymethylmethacrylate Particles Stimulate Bone Resorption of Mature Osteoclasts in  
 941 Vitro.” *Acta Orthopaedica* 79 (2): 281–88. <https://doi.org/10.1080/17453670710015166>.

942 Zhang, Q., Y. Zhao, J. Li, and H. Shi. 2020. “Microplastics in Food: Health Risks.” In .  
 943 Springer. [https://doi.org/10.1007/698\\_2020\\_453](https://doi.org/10.1007/698_2020_453).

944 Zhang, S., X. Yang, H. Gertsen, P. Peters, T. Salánki, and V. Geissen. 2018. “A Simple  
 945 Method for the Extraction and Identification of Light Density Microplastics from Soil.”  
 946 *Science of The Total Environment* 616–617 (March): 1056–65.  
 947 <https://doi.org/10.1016/J.SCITOTENV.2017.10.213>.

948 Zhang, Y., T. Gao, S. Kang, and M. Sillanpää. 2019. “Importance of Atmospheric Transport  
 949 for Microplastics Deposited in Remote Areas.” *Environmental Pollution* 254  
 950 (November): 112953. <https://doi.org/10.1016/J.ENVPOL.2019.07.121>.


951 Zhang, Y., S. Kang, S. Allen, D. Allen, T. Gao, and M. Sillanpää. 2020. “Atmospheric  
 952 Microplastics: A Review on Current Status and Perspectives.” *Earth-Science Reviews*  
 953 203 (April): 103118. <https://doi.org/10.1016/J.EARSCIREV.2020.103118>.

954 Zhang, Y., J. Lu, J. Wu, J. Wang, and Y. Luo. 2020. “Potential Risks of Microplastics

955 Combined with Superbugs: Enrichment of Antibiotic Resistant Bacteria on the Surface  
956 of Microplastics in Mariculture System.” *Ecotoxicology and Environmental Safety* 187:  
957 109852. <https://doi.org/https://doi.org/10.1016/j.ecoenv.2019.109852>.  
958 Zhu, X., L. Qiang, H. Shi, and J. Cheng. 2020. “Bioaccumulation of Microplastics and Its in  
959 Vivo Interactions with Trace Metals in Edible Oysters.” *Marine Pollution Bulletin* 154:  
960 111079. <https://doi.org/https://doi.org/10.1016/j.marpolbul.2020.111079>.  
961  
962



**Figure 1**



*Figure 1 - Flow diagram to illustrate the potential human health effects of microplastics.*

*Dotted lines represent current speculative research. (1) Giuseppe et al. 2003, (2) Peretz et al. 2014, (3) Ait Bamai et al. 2014b, (4) Gasperi et al. 2018 (5) Tang et al. 2020, (6) Valavanidis et al. 2013, (7) Prata 2018, (8) Gallagher et al. 2015, (9) Lu et al. 2016, (10) Qiao et al. 2019, (11) Li et al. 2020, (12) Morris and Acheson 2003, (13) Kirstein et al. 2016, (14) Yang et al. 2019a, (15) Lu et al. 2019.*