

**Fast versus gradual adaptation of soft monthly contact lenses in neophyte wearers**

<sup>1</sup>James S. Wolffsohn PhD [j.s.w.wolffsohn@aston.ac.uk](mailto:j.s.w.wolffsohn@aston.ac.uk)

<sup>2,3</sup>Neema Ghorbani-Mojarrad PhD [N.GhorbaniMojarrad@bradford.ac.uk](mailto:N.GhorbaniMojarrad@bradford.ac.uk)

<sup>4</sup>Marta Vianya-Estopa PhD [marta.vianya@aru.ac.uk](mailto:marta.vianya@aru.ac.uk)

<sup>5</sup>Manbir Nagra PhD [nagra.manbirk@gmail.com](mailto:nagra.manbirk@gmail.com)

<sup>6</sup>Byki Huntjens PhD [Byki.Huntjens.1@city.ac.uk](mailto:Byki.Huntjens.1@city.ac.uk)

<sup>7</sup>Louise Terry PhD [TerryL1@cardiff.ac.uk](mailto:TerryL1@cardiff.ac.uk)

<sup>8</sup>Laura Elaine Sweeney PhD [laura.sweeney@gcu.ac.uk](mailto:laura.sweeney@gcu.ac.uk)

<sup>1</sup>Debarun Dutta PhD [d.dutta@aston.ac.uk](mailto:d.dutta@aston.ac.uk)

<sup>9</sup>Mahesh Raj Joshi PhD [mahesh.joshi@plymouth.ac.uk](mailto:mahesh.joshi@plymouth.ac.uk)

<sup>1</sup>Daniel Wright BSc [yohdan1993@gmail.com](mailto:yohdan1993@gmail.com)

<sup>2</sup>Heather Bruce BSc [H.M.Bruce@bradford.ac.uk](mailto:H.M.Bruce@bradford.ac.uk)

<sup>2</sup>Emily Hallam BSc [E.Hallam@bradford.ac.uk](mailto:E.Hallam@bradford.ac.uk)

<sup>2</sup>Louise Jolly BSc [L.Jolly3@bradford.ac.uk](mailto:L.Jolly3@bradford.ac.uk)

<sup>8</sup>Ying Bing Chung BSc [YCHUNG201@caledonian.ac.uk](mailto:YCHUNG201@caledonian.ac.uk)

<sup>8</sup>Joanne Ru En Tsen BSc [RTSEN200@caledonian.ac.uk](mailto:RTSEN200@caledonian.ac.uk)

<sup>7</sup>Anya Bishop BSc [anyabishop@talktalk.net](mailto:anyabishop@talktalk.net)

<sup>7</sup>Rhona Davison BSc [davis200r@aol.com](mailto:davis200r@aol.com)

<sup>10</sup>Carole Maldonado-Codina PhD [carole.m-codina@manchester.ac.uk](mailto:carole.m-codina@manchester.ac.uk)

and members of the British and Irish University and College Contact Lens Educators (BUCCLE) group

<sup>1</sup>Optometry and Vision Science Research Group, Optometry School, Aston University, Birmingham UK

<sup>2</sup>School of Optometry and Vision Science, University of Bradford, Bradford, UK

<sup>3</sup>Wolfson Centre for Applied Health Research, Bradford Royal Infirmary, Bradford, UK

<sup>4</sup>Vision and Hearing Sciences Research Centre, Anglia Ruskin University, Cambridge, UK

<sup>5</sup>Vision and Eye Research Institute (VERI) Anglia Ruskin University, Cambridge, CB1 2LZ

<sup>6</sup>Division of Optometry and Visual Sciences, City, University of London, London, UK

<sup>7</sup>School of Optometry and Vision Sciences, Cardiff University, Cardiff, UK

<sup>8</sup>Department of Vision Sciences, Glasgow Caledonian University, Glasgow, UK

<sup>9</sup>Eye and Vision Research Group, School of Health Professions, University of Plymouth, United Kingdom

<sup>10</sup>Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK

**Corresponding Author:**

Dr Marta Vianya-Estopa

Vision and Hearing Sciences Research Centre, Anglia Ruskin University, Cambridge, CB1 1PT, UK

Tel: +44 1223 363271 Email: [marta.vianya@aru.ac.uk](mailto:marta.vianya@aru.ac.uk)

ORCID iD Marta Vianya-Estopa 0000-0001-5913-0764

**Declarations of interest:** the authors have no conflicts of interest and have no financial disclosures.

**Funding:** BUCCLE's mission is to enhance optometry education in the UK and in this pursuit is currently funded by the following companies: Alcon, Bausch + Lomb, CooperVision, Johnson and Johnson Vision, No7 Contact Lenses and Menicon. The research was designed and implemented independently. Other BUCCLE members include: Alison Alderson (University of Bradford), Arnold Cochrane (Ulster University), Claire McDonnell (TU Dublin), Dean Dunning (Bradford College), Eilidh Martin (Glasgow Caledonian University), Katharine Evans (Cardiff University), Shehzad Naroo (Aston University), Claire Mallon (The University of Manchester), Jo Underwood (Association of British Dispensing Opticians), Asma Zahidi (University of Plymouth), Orla Murphy (TU Dublin), Emma Gibson (UWE Bristol), Nas Hinglotwala (The University of Huddersfield), Marie-Laure Orr (University of Central Lancashire), Mark Mayhew (City University), Orla Murphy (TU Dublin) and Nery Garcia-Porta (Anglia Ruskin University).

## **Fast versus gradual adaptation of soft monthly contact lenses in neophyte wearers**

### **Abstract**

**Aim:** To determine if a gradual adaptation period is necessary for neophytes when fitted with modern hydrogel or silicone hydrogel reusable disposable contact lenses.

**Method:** Across four sites, 74 neophytes (18-28 years) were randomly assigned to a reusable lens: Proclear® (hydrogel) or Biofinity® (silicone hydrogel) and an adaptation schedule: fast (10 hours wear from the first day) or gradual (4 hours on the first day, increasing their wear time by 2 hours on each subsequent day until they had reached 10 hours). Masked investigators graded ocular surface physiology and non-invasive tear breakup time (NIBUT) and a range of comfort, vision and lens handling subjective ratings (0-100 visual analogue scales) were recorded at the baseline visit and after 10 hours of lens wear, 4-6 days and 12-14 days after lens fitting. Subjective scores were also repeated after 7 days.

**Results:** There was no difference ( $p>0.05$ ) in ocular surface physiology or NIBUT between fast and gradual adaptation groups at any time point in either lens type with the exception of increased corneal staining ( $p=0.019$ ) in the silicone hydrogel fast adaptation group after 4-6 days. Subjective scores were also similar across the visits and lens types with the exception of 'lens awareness' ( $p=0.019$ ) which was less in the gradual versus the fast adaptation silicone hydrogel lens group at 12-14 days.

**Conclusion:** There seems to be no clinical benefit for recommending a gradual adaptation period in new wearers fitted with modern soft reusable disposable contact lenses. The findings of this work add to a growing body of evidence suggesting that such advice is unnecessary in regular soft contact lens wear, which has important ramifications for the initial clinical management of these patients.

**Key words:** Soft contact lens, reusable, adaptation, neophyte, fast, gradual

## 1. Introduction

Currently, conventional practice advocates a cautious 'easing-in' approach for adapting new contact lens wearers (neophytes)[1]. In daily lens wear, this usually involves wear schedules of 2 to 4 hours on the first day followed by increases of 1-2 hours daily until the desired wear time is achieved. Whilst this is likely to be beneficial for newly-adapting rigid lens wearers, it is less likely to be important for wearers of soft contact lenses. Soft lenses have a much lower modulus than rigid lenses [2, 3] and have less interaction with the upper eyelid due to a larger diameter and reduced lens movement, which makes them significantly more comfortable from the very first application. For this reason, many patients use soft lenses on an occasional basis and the concept of building-up of wear time in the traditional sense seems redundant under these circumstances.

Previous work from this group[4] comparing fast to gradual adaptation in neophyte daily disposable lens wearers showed no significant differences in ocular physiology over the first two weeks of lens wear. Limbal, bulbar and palpebral conjunctival redness as well as corneal staining were found to be similar for the two groups with both contemporary hydrogel and silicone hydrogel daily disposable lenses. This finding lends weight to the hypothesis that the oxygen transmissibility of a lens is not relevant in deciding if a gradual adaptation period is required in a soft lens. Furthermore, the work showed that subjective comfort, vision and lens handling were not negatively impacted by a fast adaptation schedule; in fact, lens awareness and ease of lens removal were *improved* in the fast compared to the gradual adapters in the hydrogel lens wearers.

The report was the first to provide evidence that eye care practitioners could eliminate gradual adaptation periods in soft lenses – at least for daily disposable wearers. However, it remains unknown whether the same principle can be applied to reusable daily wear soft contact lenses which remain the most widely prescribed lens category across the world, currently making-up up 44% of lens fits globally [5]. There are additional complexities which could influence comfort and adaptation with reusable lenses compared with daily disposable lenses, such as the interaction of the care regimen with the ocular surface [6, 7] as well as the potential for increased levels of deposition and its effect on ocular physiology [8].

This work set out to build upon the findings of previous work[4] and sought to gain a better understanding of whether the recommendation of gradual adaptation was supported for reusable daily wear hydrogel and silicone hydrogel contact lenses. Specifically, the work aimed to investigate if there were differences in ocular surface physiology and subjective performance in contact lens neophytes prescribed reusable lenses who underwent a fast versus a gradual adaptation schedule in the first two weeks of lens wear.

## **2. Methods**

### **2.1 Study lenses and care regimen**

The two monthly reusable lenses investigated in this work were Proclear® and Biofinity® (CooperVision Inc.) (Table 1). These lenses were selected based on the similarity of their design (e.g. lens edge shape) and as representative examples of commonly prescribed hydrogel and silicone hydrogel monthly reusable lenses. Participants were fitted with one of the two lens types and worn bilaterally (as a matching lens pair) on a daily wear, reusable basis for a period of 12-14 days.

All participants used Opti-Free® Puremoist® multi-purpose contact lens solution (Alcon Laboratories Inc.) throughout the study together with the manufacturer-provided flat lens case. The care regimen is described as a buffered solution containing the dual disinfectants/preservatives POLYQUAD® (polyquartanium-1) 0.001% and ALDOX® (myristamidopropyl dimethylamine) 0.0006%[9] . Two wetting agents; Tetronic 1304 (BASF Corporation) and a proprietary linear diblock copolymer composed of poly(oxyethylene)-poly-(oxybutylene) named EOBO, HydraGlyde® Moisture Matrix are present as well as sodium citrate, sodium chloride, boric acid, sorbitol, aminomethylpropanol and disodium EDTA. Participants were instructed to use the solution following the manufacturer guidelines which also included a rub-and-rinse step.

### **2.2 Study Design**

This was a prospective, parallel-group, randomised, investigator-masked, multi-site study based at four academic institutions: Aston University (Birmingham, UK), University of Bradford (Bradford, UK), Cardiff University (Cardiff, UK), and Glasgow

Caledonian University (Glasgow, UK). All four sites received human ethics approval from their respective institutional research ethics committee. The study conformed to the tenets of the Declaration of Helsinki and all participants provided written informed consent prior to enrolment.

Inclusion criteria included being aged between 18-40 years with astigmatism  $\leq 0.75$ DC, being deemed suitable for contact lens wear following anterior eye assessment and being in possession of an in-date spectacle prescription. Participants were excluded if they had a history of contact lens wear within the previous six months, were pregnant or breast-feeding, had had recent refractive surgery, had a known hypersensitivity to saline or sodium fluorescein, took medications known to affect contact lens wear or had a systemic or ocular condition that could affect lens wear. The sample size of participants required for the study was estimated using power calculations from a previous study using daily disposable lenses[4]: 10 participants in each adaptation/lens material group would have 80% power to detect a difference of at least 10 points on a 0-100 grading scale for subjective scores.

Participants attended three visits. At the initial visit, baseline investigations included refraction, visual acuity, non-invasive tear breakup time (NIBUT) using either a Tearscope Plus (Keeler, Windsor, UK) or keratometry mires (Bausch and Lomb Rochester NY, USA) and slit lamp examination of the ocular surface: bulbar, limbal and palpebral conjunctival hyperaemia, palpebral roughness, and corneal staining were graded to the nearest 0.1 units using Efron grading scales[10]. The assessment was performed using 16x magnification under white light with the addition of sodium fluorescein (1.5 mg impregnated strips) for the observation of corneal staining using blue light together with a yellow enhancement filter in front of the observation system[11].

All eligible participants at each site were assigned to one of the two lenses for the investigation, with each site only fitting one of the lens types. Lens fit was assessed using the simplified approach proposed by Wolffsohn and colleagues[12]. An unacceptable fit was identified by the presence of limbal excursion or if there were two or more minus grading values for the fitting parameters. Subjective responses were reported using 0-100 visual analogue scales where 0 indicated a very poor or negative

experience and 100 indicated a very positive experience. At initial lens dispensing the following were recorded: 'comfort before lens application', 'overall comfort' and 'visual quality'.

Participants were then randomly allocated to one of two adaptation schedules; i) no build-up of wearing time (fast adaptation) where participants would wear lenses for 10 hours from the first day or ii) a more gradual build-up (gradual adaptation) where participants would wear lenses for 4 hours on the first day and increase their wear time by 2 hours on each subsequent each day until they reached 10 hours. Investigators collecting data were masked to the adaptation schedule group. All participants were instructed fully on contact lens application and removal and given full instructions on how to care for their lenses, including the use of the care regimen.

Participants returned to the clinic for two further follow-up visits once they had reached 10 hours of lens wear: i) 4-6 days and ii) 12-14 days after fitting. Slit lamp biomicroscopy and NIBUT assessments were carried out at both visits similarly to the initial baseline visit. The following subjective scores were recorded using 0-100 visual analogue scales: 'comfort prior to lens application', 'overall comfort', 'vision quality', 'lens awareness throughout the day', 'end-of-day comfort', 'ease of lens application' and 'ease of lens removal'. Participants were also asked to record these same parameters after wearing the lenses for 7 days and to return the completed questionnaire at the final visit.

### **2.3 Statistical Analysis**

Statistical analyses were performed using IBM SPSS Statistics (v23 IBM Corp. Chicago, Illinois, USA). The data were not found to be normally distributed (Kolmogorov-Smirnov Test  $p < 0.05$ ) therefore Mann-Whitney U tests were used to investigate the differences between the gradual and fast adaptation groups at each visit. The statistical significance level was set at  $p < 0.05$ .

### 3. Results

Seventy-four participants were enrolled and the demographics of the study groups are shown in Table 2. Overall the age range of all the study cohorts remained similar between 18-28 years, and the range of refractive error (spherical equivalent) was between +0.25 and -6.50 DS. All recruited participants completed the study and no adverse events occurred. No lens fits were deemed 'unacceptable'.

#### 3.1 Ocular surface physiology and tear film stability

There were no statistically significant differences ( $p>0.05$ ) in ocular surface physiology or NIBUT measurements between the two adaptation schedule groups at baseline, or at the two follow-up visits for the hydrogel or silicone-hydrogel wearers (Tables 3 and 4); the only exception was after 4-6 days of wear, when the gradual adaptation silicone hydrogel wearers demonstrated significantly lower scores for corneal staining compared to the fast adaptation group ( $p=0.019$ ; Table 4), but this difference was not sustained after 12-14 days of lens wear.

#### 3.2 Subjective assessments

At baseline there were no statistically significant differences ( $p>0.05$ ) in subjective scores between the two adaptation schedule groups for both the hydrogel (Table 5) and silicone hydrogel (Table 6) wearers. This was also true at 4-6 days and day 7 after lens wear commenced. After 12-14 days of silicone hydrogel lens wear, 'lens awareness' ( $p=0.02$ ) was significantly better in the gradual compared to the fast adaptors, but there were no other differences between the adaptation schedules (Table 5 and Table 6).

### 4. Discussion

This study built upon the knowledge gained from the first investigation on this topic which compared the effect of a fast compared to a more traditional gradual adaptation schedule on ocular surface physiology and subjective acceptance in neophyte daily disposable lens wearers[4]. As far as possible the same methodology and statistical analyses were repeated for the current second sister study, this time, using reusable daily wear contact lenses. Overall, the results from the present work are similar to those found in the previous study. Neither a fast nor a gradual adaptation schedule



had any major impact on the short-term ocular surface physiology or tear film stability with modern hydrogel or silicone hydrogel reusable contact lenses.

In hydrogel lens wear there were no differences between adaptation groups for bulbar, limbal or palpebral hyperaemia across time points and this was also the case for palpebral roughness and corneal staining. Similar results were seen in the silicone hydrogel lens wearers except that the gradual adaptation group demonstrated reduced levels of corneal staining after 4-6 days compared with the fast group. Given that the corneal staining scores were 0.3 versus 0.1 Efron grading units (fast vs. gradual adaptation groups, respectively), it seems reasonable to conclude that these differences are not clinically significant since their magnitude lies within the 'normal' range on this grading scale[13]. Any differences between the two groups in this parameter had disappeared by 12-14 days.

Contact lens wear causes disruption to the normal tear film structure and function [14-16] which is thought to be a significant factor in negatively impacting ocular discomfort despite the lack of conclusive evidence linking the two. No differences were observed between the two adaptation schedules for NIBUT in either lens type at either visit which suggests that tear film stability is not adversely affected as a result of how quickly the wearing time is built up in reusable lenses. Overall, these results are very similar to earlier findings investigating adaptation schedule in daily disposable wearers[4] with the exception that in the daily disposable work a longer NIBUT was found in those undergoing a gradual adaptation in silicone hydrogel lenses at the 12-14 day visit.

In terms of subjective comfort-related responses, there were no statistically significant differences between the two adaptation groups in the hydrogel lens wearers. Interestingly, in hydrogel daily disposable wearers 'lens awareness' and 'end-of-day' comfort were shown to be better in the fast versus the gradual adaptation group after 7 and 12-14 days, respectively[4]. No such differences have been demonstrated in the current work which could be as a result of the particular hydrogel lens design chosen, lens deposition differences or factors related to the lens/solution combination. This lack of comfort-related symptoms difference between the two adaptation groups is in

line with no differences being observed in ocular physiology and NIBUT in this lens type.

In the silicone hydrogel wearers, 'lens awareness' scores were better (i.e. scores were higher which corresponded to reduced lens awareness) in the gradual adaptation versus the fast adaptation group at the 12-14 day visit and this difference (86 vs. 71) is quite marked. The gradual adaptors also presented with significantly reduced corneal staining at the 4-6 day time point and it is not clear if this could have contributed to the subsequent lens awareness increases in this group at the following visit. Previous work has shown a link between comfort and levels of SICS staining[17-19], yet it is unlikely that the use of other lens care solutions such as hydrogen peroxide would have reduced the level of corneal staining observed or changed the study outcome as the frequency of cleaning was the same between the fast and gradual adaptation groups. It would be interesting to investigate whether or not this 'lens awareness' difference persists longer-term, but the difference between the two adaptation groups in this lens type is somewhat offset by there being no other differences in subjective comfort scores over the two-week study period.

Visual quality was similar for the adaptation groups in both lens types at all time points across the two-week period, which is in line with previous findings for daily disposable lenses. This study also evaluated subjective handling aspects relating to 'ease of application' and 'ease of removal' at each follow-up visit; as with the daily disposable lens study, no significant differences were found between the fast and gradual adaptation groups at any of the time points or for either lens type. This result is not unexpected given that the total amount of handling time is the same whichever adaptation schedule is followed i.e. participants would be applying and removing the lenses once per day.

Overall, the results from this work suggest that gradual adaptation to modern spherical reusable disposable soft lenses is unnecessary, regardless of the oxygen permeability of the material. As has been previously stated, this does not mean that wearers should be instructed to wear their lenses for 10 hours from the start regardless, but rather a sensible approach would be to instruct patients to wear them for as long as they are comfortable up to a suggested maximum. The first few weeks of lens wear are very

important in terms of the long-term success of a new contact lens wearer so the patient should be followed up to determine whether they have any issues that need addressing[20].

This work with reusable soft contact lenses has added to the growing body of evidence showing that gradual adaptation in neophytes has little clinical benefit compared to a fast adaptation approach in both hydrogel and silicone hydrogel lenses. These findings have important ramifications for the clinical management of these patients in the initial lens wear period.

## References

- [1] Morgan P. Optimising contact lens wear for a lifetime of use. *Optician* 2013;245(6400).
- [2] Bhamra TS, Tighe BJ. Mechanical properties of contact lenses: The contribution of measurement techniques and clinical feedback to 50 years of materials development. *Cont Lens Anterior Eye* 2017;40(2):70-81.
- [3] Carney LG, Mainstone JC, Carkeet A, Quinn TG, Hill RM. Rigid lens dynamics: lid effects. *The CLAO journal : official publication of the Contact Lens Association of Ophthalmologists, Inc* 1997;23(1):69-77.
- [4] Wolffsohn JS, Dhirajlal H, Vianya-Estopa M, Nagra M, Madden L, Sweeney LE, et al. Fast versus gradual adaptation of soft daily disposable contact lenses in neophyte wearers. *Cont Lens Anterior Eye* 2020;43(3):268-73.
- [5] Morgan P, Woods CA, Tranoudis, Efron N, Merchan L, Grupcheva., et al. International Contact Lens Prescribing in 2020. 35. *Contact Lens Spectrun* 2020:26-32.
- [6] Carnt NA, Evans VE, Naduvilath TJ, Willcox MD, Papas EB, Frick KD, et al. Contact lens-related adverse events and the silicone hydrogel lenses and daily wear care system used. *Arch Ophthalmol* 2009;127(12):1616-23.
- [7] Lazon de la Jara P, Papas E, Diec J, Naduvilath T, Willcox MD, Holden BA. Effect of lens care systems on the clinical performance of a contact lens. *Optom Vis Sci* 2013;90(4):344-50.
- [8] Zhao Z, Naduvilath T, Flanagan JL, Carnt NA, Wei X, Diec J, et al. Contact lens deposits, adverse responses, and clinical ocular surface parameters. *Optom Vis Sci* 2010;87(9):669-74.
- [9] Campbell R, Kame G, Leach N, Paul M, White E, Zigler L. Clinical benefits of a new multipurpose disinfecting solution in silicone hydrogel and soft contact lens users. *Eye Contact Lens* 2012;38(2):93-101.
- [10] Efron N. Grading scales for contact lens complications. *Ophthalmic Physiol Opt* 1998;18(2):182-6.
- [11] Peterson RC, Wolffsohn JS, Fowler CW. Optimization of anterior eye fluorescein viewing. *American journal of ophthalmology* 2006;142(4):572-5.
- [12] Wolffsohn JS, Hunt OA, Basra AK. Simplified recording of soft contact lens fit. *Cont Lens Anterior Eye* 2009;32(1):37-42.
- [13] Efron N, Morgan PB, Katsara SS. Validation of grading scales for contact lens complications. *Ophthalmic Physiol Opt* 2001;21(1):17-29.
- [14] Nichols JJ, Mitchell GL, King-Smith PE. Thinning rate of the precorneal and prelens tear films. *Invest Ophthalmol Vis Sci* 2005;46(7):2353-61.
- [15] Glasson MJ, Stapleton F, Keay L, Willcox MD. The effect of short term contact lens wear on the tear film and ocular surface characteristics of tolerant and intolerant wearers. *Cont Lens Anterior Eye* 2006;29(1):41-7; quiz 9.
- [16] Nichols JJ, King-Smith PE. The impact of hydrogel lens settling on the thickness of the tears and contact lens. *Invest Ophthalmol Vis Sci* 2004;45(8):2549-54.
- [17] Diec J, Evans VE, Tilia D, Naduvilath T, Holden BA, Lazon de la Jara P. Comparison of ocular comfort, vision, and SICS during silicone hydrogel contact lens daily wear. *Eye Contact Lens* 2012;38(1):2-6.
- [18] Dutta D, Sarkes M, Suneetha N, Markoulli M. Repeatability and correlation of tear ferning with lipid layer thickness, non-invasive tear break up time and OSDI. *American Academy of Optometry*. Anaheim; 2016.
- [19] Kumar M, Shetty R, Jayadev C, Rao HL, Dutta D. Repeatability and agreement of five imaging systems for measuring anterior segment parameters in healthy eyes. *Indian J Ophthalmol* 2017;65(4):288-94.
- [20] Sulley A, Young G, Hunt C. Factors in the success of new contact lens wearers. *Cont Lens Anterior Eye* 2017;40(1):15-24.

403 [21] Kerr C, Chauhan K. The ACLM contact lens year book 2020.: Devizes: Association  
404 of Contact Lens Manufacturers; 2020.

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

Lens name	Biofinity®	Proclear®
Manufacturer	CooperVision Inc.	CooperVision Inc.
Material	Comfilcon A	Omafilcon B
Base Curve (mm)	8.6	8.6
Total Diameter (mm)	14.0	14.2
Water content (%)	48	62
Oxygen permeability (ISO units)	96	20
Back vertex power range (BVP)	+8.00 to -12.00D	+6.50 to -20.00D

436

437 **Table 1:** Study lenses (parameters from the ACLM Yearbook)[21]

438

439

<b>Lens</b>	<b>Experimental Group</b>	<b>Participants</b>	<b>Age (years)</b>	<b>Male/Female Ratio</b>	<b>Refraction (Spherical equivalent in dioptries)</b>
Biofinity®	Gradual	17	18 - 23	4 / 13	+0.25 to -6.50D
	Fast	18	18 - 28	1 / 17	+0.25 to -4.50D
Proclear®	Gradual	20	18 - 27	3 / 17	+0.50 to -5.25D
	Fast	19	18 - 28	6 / 13	+0.50 to -6.50D

440

441 **Table 2:** Demographic and refractive details of the study participants.

		Baseline			Day 4-6			Day 12-14		
		Mean	SD	P	Mean	SD	P	Mean	SD	P
Bulbar Hyperaemia	Fast	0.8	±0.6	0.879	0.7	±0.7	0.365	1.0	±0.7	0.351
	Gradual	0.8	±0.8		0.9	±0.6		0.7	±0.8	
Limbal Hyperaemia	Fast	0.6	±0.4	0.945	0.8	±0.4	0.322	0.7	±0.6	0.322
	Gradual	0.6	±0.6		0.6	±0.6		0.5	±0.6	
Palpebral Hyperaemia	Fast	0.8	±0.5	0.728	0.8	±0.7	0.771	0.8	±0.6	0.513
	Gradual	0.7	±0.6		0.8	±0.6		0.7	±0.7	
Palpebral Roughness	Fast	0.7	±0.5	0.444	0.6	±0.6	0.879	0.6	±0.7	0.513
	Gradual	0.6	±0.5		0.6	±0.5		0.4	±0.5	
Corneal Staining	Fast	0.3	±0.4	0.771	0.5	±0.5	0.559	0.5	±0.4	0.708
	Gradual	0.1	±0.2		0.3	±0.4		0.5	±0.4	
Non-invasive breakup time (s)	Fast	9.1	±1.4	0.999	8.2	±1.1	0.270	7.6	±1.3	0.351
	Gradual	9.0	±1.6		8.7	±1.2		8.1	±1.6	

**Table 3:** Comparison of ocular physiology in fast and gradual adaptation of neophytes fitted with reusable hydrogel soft contact lenses. Efron scale grading between 0 and 4 units, using 0.1 increments. SD = standard deviation; p = significance value. (bold indicates level <0.05).



		Baseline			Day 4-6			Day 12-14		
		Mean	SD	P	Mean	SD	P	Mean	SD	P
Bulbar Hyperaemia	Fast	0.6	±0.4	0.335	0.8	±0.4	0.173	0.9	±0.4	0.883
	Gradual	0.7	±0.4		0.8	±0.4		0.9	±0.4	
Limbal Hyperaemia	Fast	0.5	±0.4	0.636	0.6	±0.4	0.883	0.7	±0.3	0.660
	Gradual	0.5	±0.4		0.5	±0.3		0.8	±0.4	
Palpebral Hyperaemia	Fast	0.4	±0.3	0.590	0.4	±0.4	0.393	0.5	±0.5	0.405
	Gradual	0.5	±0.4		0.5	±0.4		0.6	±0.5	
Palpebral Roughness	Fast	0.4	±0.2	0.732	0.4	±0.2	0.463	0.5	±0.3	0.935
	Gradual	0.3	±0.2		0.5	±0.4		0.5	±0.4	
Corneal Staining	Fast	0.2	±0.3	0.999	0.3	±0.3	<b>0.019</b>	0.3	±0.4	0.351
	Gradual	0.1	±0.2		0.1	±0.2		0.2	±0.3	
Non-invasive breakup time (s)	Fast	11.1	±3.2	0.590	10.1	±2.0	0.270	9.6	±2.9	0.613
	Gradual	11.1	±2.3		10.4	±2.6		10.1	±4.2	

**Table 4:** Comparison of ocular physiology in fast and gradual adaptation of neophytes fitted with reusable silicone hydrogel soft contact lenses. Efron scale grading between 0 and 4 units, using 0.1 increments. SD = standard deviation; p = significance value. (bold indicates level <0.05).

457

458

		Baseline			Day 4-6			Day 7			Day 12-14		
		Me	SD	p	Me	SD	p	Me	SD	p	Me	SD	p
Comfort prior to lens wear	Fast	97.9	±4.2	0.4	96.1	±7.6	0.41	97.1	±5.6	0.92	98.7	±3.3	0.74
	Gradual	99.0	±4.5	28	97.5	±7.3	1	96.8	±7.8	3	96.3	±8.7	9
Overall comfort	Fast	85.5	±9.1	0.7	81.1	±15.4	0.46	78.7	±17.5	0.47	82.4	±13.3	0.35
	Gradual	86.0	±11.4	08	85.0	±13.2	1	83.0	±16.2	8	86.0	±12.3	1
Visual quality	Fast	93.2	±8.0	0.6	94.2	±7.9	0.12	91.6	±10.5	0.33	95.3	±6.3	0.28
	Gradual	91.5	±15.2	87	85.5	±17.5	7	86.5	±14.5	6	88.8	±16.7	3
Lens Awareness	Fast				77.9	±16.3	0.62	76.7	±18.8	0.47	76.6	±22.1	0.51
	Gradual				75.0	±18.9	7	73.6	±17.4	8	74.5	±18.3	3
End of Day Comfort	Fast				72.9	±22.6	0.83	70.3	±24.4	0.94	70.8	±22.7	0.96
	Gradual				75.0	±22.9	5	73.5	±20.7		72.8	±17.3	7
Ease Application	Fast				84.7	±12.9	0.84	85.8	±15.2	0.98	88.9	±14.4	0.68
	Gradual				90.5	±11.0	6	87.8	±10.2	9	88.3	±12.2	7
Ease Removal	Fast				89.7	±11.0	0.27	92.6	±11.9	0.96	91.8	±10.3	0.56
	Gradual				94.3	±8.5	0	92.5	±12.0	7	94.8	±7.7	9

459

460

461

462

463

464

**Table 5:** Comparison of subjective ratings in fast and gradual adaptation of neophytes fitted with reusable hydrogel soft contact lenses using visual analogue scales (0-100). SD = standard deviation; p = significance value (bold indicates level <0.05).

465

466

		Baseline			Day 4-6			Day 7			Day 12-14		
		Me	SD	p	Me	SD	p	Me	SD	p	Me	SD	p
Comfort prior to lens wear	Fast	92.0	±9.9	0.6	88.9	±13.7	0.83	93.1	±9.0	0.96	93.6	±7.7	0.5
	Gradual	93.7	±9.3	60	91.7	±9.0	2	93.2	±8.8	1	95.3	±7.3	46
Overall comfort	Fast	77.5	±20.9	0.6	74.4	±21.4	0.52	81.5	±15.8	0.83	79.8	±17.9	0.3
	Gradual	80.8	±20.4	13	80.6	±13.8	5	84.6	±9.8	2	87.5	±10.0	18
Visual quality	Fast	88.8	±11.0	0.5	79.4	±18.8	0.50	87.9	±6.5	0.90	89.1	±9.1	0.7
	Gradual	83.2	±17.4	46	82.2	±20.0	3	84.6	±16.3	9	88.3	±13.4	08
Lens Awareness	Fast				74.7	±28.1	0.61	78.0	±21.7	0.73	71.0	±21.3	<b>0.0</b>
	Gradual				74.7	±19.5	8	76.1	±22.1	2	86.1	±11.5	<b>19</b>
End of Day Comfort	Fast				70.6	±26.4	0.98	73.4	±20.5	0.96	76.1	±17.1	0.3
	Gradual				74.3	±18.1	7	72.7	±19.3	1	81.3	±17.3	69
Ease Application	Fast				81.3	±15.2	0.16	82.9	±14.9	0.99	90.3	±9.8	0.7
	Gradual				69.6	±21.5	3	81.8	±15.9	9	87.4	±13.2	32
Ease Removal	Fast				83.0	±17.4	0.52	87.7	±13.4	0.93	93.6	±8.8	0.6
	Gradual				80.7	±15.3	5	87.5	±13.6	5	90.4	±13.3	13

467

468 **Table 6:** Comparison of fast and gradual adaptation of neophytes fitted with  
469 reusable silicone hydrogel soft contact lenses using visual analogue scales (0-100).  
470 SD = standard deviation; p = significance value. (bold indicates level <0.05).